

Sensory evaluation tools stroke described in Portuguese: a systematic review

Instrumentos de avaliação sensorial pós-acidente vascular encefálico (AVE) descritos em português: uma revisão sistemática

Los instrumentos de evaluación sensorial posaccidente vascular encefalico (AVE) descritos en portugués: una revisión sistemática

Jennifer Souza Fagundes¹, Aline Cristiane Binda¹, Julia Garzim Faria¹, Daniele Peres², Stella Maris Michaelsen³

ABSTRACT | Most survivors of stroke present, among other consequences, a sensory deficit. To assess the impact of this deficit on the activity and the effect of sensory reeducation protocols, it is important to use objective instruments. The objective of this systematic review was to analyze which sensory evaluation tools for post-stroke patients are available in Brazilian Portuguese and describe its features and/or properties. A search was conducted in the electronic databases SCIELO, LILACS, PUBMED, CINAHL, MEDLINE via OVDI, and EMBASE. The relevant studies were analyzed as to accessibility, objectivity of the score and measurement properties, adding articles through manual search when necessary. The search resulted in 96 studies, reduced to five eligible ones. A study was added through the list of references and the manual search was used for complementation. Among the seven reviewed articles, only three sensory evaluation tools are available in Portuguese: moving touch pressure (MTP), sensitivity domain of Fugl-Meyer Scale (FMS), and Nottingham Sensory Assessment (NSA). In addition, although the reliability of the FMS and of the NSA may be considered appropriate, the other properties of measurement need to be evaluated in future studies.

Keywords | Stroke; Sensory Modalities; Touch; Proprioception.

RESUMO | Grande parte dos sobreviventes de acidente vascular encefálico (AVE) apresenta, além de outras sequelas, algum déficit sensorial. Para avaliar o impacto desse déficit na atividade e o efeito de protocolos de reeducação sensorial é importante utilizar instrumentos objetivos. Os objetivos desta revisão sistemática foram analisar quais instrumentos de avaliação sensorial para pacientes pós-AVE estão disponíveis em português brasileiro e descrever suas características e/ou propriedades. Realizou-se uma busca nas bases de dados eletrônicas SciELO, LILACS, PubMed, CINAHL, MEDLINE via OVDI e Embase. Os estudos relevantes foram analisados quanto à acessibilidade, objetividade da pontuação e propriedades de medida, acrescentando-se artigos por meio de busca manual quando necessário. As buscas resultaram em 96 estudos, reduzidos a cinco estudos elegíveis. Um estudo foi adicionado por meio da lista de referências, e a busca manual foi utilizada para complementação. Entre os sete artigos analisados, há somente três instrumentos de avaliação sensorial disponíveis em português: Moving touch pressure (MTP), domínio sensibilidade da Escala de Fugl-Meyer (EFM) e Avaliação Sensorial de Nottingham (ASN). Além disso, embora a confiabilidade da EFM e da ASN possa ser considerada adequada, as demais propriedades de medida necessitam ser avaliadas em futuros estudos.

Study developed in the Motor Control Laboratory - LADECOM, Health and Sport Science Center - CEFID, Universidade do Estado de Santa Catarina - UDESC - Florianópolis (SC), Brazil.

¹Physical therapist, Universidade Estadual do Centro Oeste; graduate in Neurofunctional Rehabilitation: Advances of Neuroscience for Rehabilitation, Faculdade Inspirar - Curitiba (PR), Brazil.

²MSc, Master in Physiotherapy, Universidade do Estado de Santa Catarina CEFID/UDESC - Florianópolis (SC), Brazil.

³PhD, Associate Professor, Department of Physiotherapy, doctor in Rehabilitation; Graduate Program in Physiotherapy, Universidade do Estado de Santa Catarina CEFID/UDESC - Florianópolis (SC), Brazil.

Descritores | Acidente Vascular Cerebral; Modalidades Sensoriais; Tato; Propriocepção.

RESUMEN | Gran parte de los supervivientes de accidente vascular encefálico (AVE) presentan, además de otras secuelas, alguno déficit sensorial. Para evaluar el impacto de ese déficit en la actividad y el efecto de protocolos de reeducación sensorial es importante utilizar instrumentos objetivos. Los objetivos de esta revisión sistemática son analizar cuales instrumentos de evaluación sensorial para pacientes pos-AVE están disponibles en portugués brasileño y describir sus características y/o propiedades. Se realizó una búsqueda en las bases de datos electrónicos SciELO, LILACS, PubMed, CINAHL, MEDLINE vía OVDI y Embase. Los estudios relevantes fueron analizados en relación a la accesibilidad, objetividad de la puntuación y

propiedades de medida, añadiéndose artículos mediante búsqueda manual cuando necesario. Las búsquedas resultaron en 96 estudios, reducidos a cinco estudios elegibles. Se agregó un estudio mediante la lista de referencias, y se utilizó la búsqueda manual para la complementación. Entre los siete artículos analizados, hay solamente tres instrumentos de evaluación sensorial disponibles en portugués: Moving touch pressure (MTP), dominio sensibilidad de la Escala de Fugl-Meyer (EFM) y Evaluación Sensorial de Nottingham (ESN). Además, aunque la confiabilidad de la EFM y de la ESN es considerada adecuada, las demás propiedades de medida necesitan ser evaluadas en estudios futuros.

Palabras clave | Acidente Vascular Cerebral; Modalidades Sensoriales; Tato; Propriocepção.

INTRODUCTION

Stroke are the leading cause of death in the adult population of Brasil¹ and one of the main reasons of disability in the world, since 70% of patients do not return to a productive life². The classic sign caused by stroke is hemiplegia or hemiparesis, however most survivors present some sensorial deficit^{3,4} that varies in intensity, area, and mode, and that is not well rated in routine exams^{3,5}.

In the acute phase after stroke, sensory changes may reach 85% of cases⁵. For Tyson et al.⁴ and Wagner et al.¹², the impairment of the tactile sensitivity is more common than the proprioceptive one, while Conell, Lincoln and Radford³ argue that proprioception and stereognosis are more often impaired. Scalha et al.⁷ evaluated the sensitivity in the affected upper limb of 20 patients with hemiparesis. All showed impairment in at least one of the sensory modalities evaluated, and, respectively, only 5% and 20% of the sample presented tactile discrimination or preserved stereognosis.

Sensory function is related to the motor function^{6,7}, in a way that sensory deficits interfere in the performance of activities and in the rehabilitation⁷. Individuals without sensory changes present better motor recovery^{8,9}, better levels of activity and a reduction in the length of hospital stay¹⁰. However, the sensory recovery, after stroke, receives less attention than motor recovery¹¹.

Sensitivity is related to mobility and independence in daily life activities⁴. Deficits in ankle proprioception change the walking intensity^{13,14} and interfere in the distance walked in the Six-Minute Walk Test¹⁵.

The knowledge of sensory deficits helps choosing rehabilitation strategies¹⁶. A systematic review¹⁷ found 16 sensory measuring tools, but 11 were not available for use, with few measures undertaken to provide consistent data and easy application. There is a need for standardization, and specific and reliable tests to evaluate the sensory condition of post-stroke patients¹⁸. It is necessary to determine the most relevant focus to recover the somatosensory functions⁸ creating sensory rehabilitation protocols in different stages of recovery¹⁹.

In this context, the objectives of this study were (1) to review systematically the standardized sensory assessment tools and with quantitative results available in Portuguese to evaluate patients affected by stroke; and (2) to describe their features and/or measurement properties.

METHODOLOGY

This systematic review followed the recommendations of PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) when pertinent²⁰.

The search aimed to identify evaluation instruments translated and/or validated to Portuguese. Articles published until March 2013 were searched in the databases: MEDLINE via OVDI, CINAHL, PUBMED, EMBASE – keywords and operators were (sensation OR touch OR tactile alteration OR proprioception OR joint position OR joint movement) AND (scales OR measures OR instruments OR clinical assessment tools) AND (stroke OR hemiplegia

OR hemiparesis) AND (portuguese OR brazil OR brazilian); LILACS and SCIELO – (sensação OR toque OR alteração tátil OR propriocepção OR posição articular OR movimento articular) AND (escalas OR mensurações OR instrumentos OR instrumentos de avaliação clínica) AND (Acidente Vascular Encefálico OR hemiplegia OR hemiparesia) AND (português OR Brasil OR brasileiro). The titles/abstracts were checked by two researchers to select the relevant articles.

For selection, the articles were reviewed independently and blindly by two other researchers, according to the following inclusion criteria: *Characteristic* – study that evaluates sensitivity; *Participants* – adults >18 years diagnosed with stroke; *Accessibility* – form of application described in Brazilian Portuguese, available online and/or by contacting the author*; *Test* – tactile sensitivity and/or proprioceptive sensitivity and/or stereognosis; *Measure* – test that results in measure/objective outcome; *Reliability* – described in Portuguese or in the original version. *If the translation process has generated an article in English, we contacted the original authors to check the availability of the instrument and the description of the application in Portuguese. We only included the instrument in the study when their availability was identified. The exclusion criterion was the non-standardization of the test application.

Microsoft Excel® spreadsheets with the checklist of inclusion criteria of each evaluator were compared and, when there were disagreements, the final inclusion was discussed with a third author. Theses/dissertations were consulted for completion when found in manual search, which was carried out to identify the description of the test's application in Portuguese.

For data extraction, we analyzed studies based on the description of the test in Brazilian Portuguese, evaluated sensory mode, form of application, score and measurement properties²¹, when described. The studies were evaluated according to the Consensus-based Standards for the selection of health Measurement Instruments (COSMIN)²². The COSMIN checklist follows three steps and is recommended for use in systematic reviews of measurement properties. Step one of COSMIN is the verification of which measurement properties were evaluated (Internal Consistency, Reliability, Measurement Error, Content Validity, Structural Validity Test of Hypotheses, Cross-cultural Validation, Criterion Validity and Responsiveness).

Step two is to determine if the statistical method used in the article is based on the classical test theory (CTT) or on the item response theory (IRT). Step three is to determine if a study complies with the requirements of good methodological quality and constitutes in boxes of evaluation for several specific items that are classified as excellent, good, reasonable, and poor.

RESULTS

Searches resulted in 96 studies, 10 evaluated by the inclusion/exclusion criteria and the rest deleted by the reading of title/abstract or if they were duplicates. Of ten reviewed articles, five did not present the inclusion criteria and the other five described the application of three instruments for sensitivity evaluation in post-stroke individuals: Moving touch pressure^{23,24}, sensitivity domain of the Fugl-Meyer Scale^{7,25}, and Nottingham Sensory Assessment^{7,19}. Of the studies above, three were published in English^{7,23,24}. By manual search, we identified the full description of a test application²⁶ and, in the list of references, we identified another study concerning the Fugl-Meyer Scale²⁷ (Figure 1).




Figure 1. Flowchart of the systematic review

Two instruments were validated for the Brazilian population and had their reliability assessed showing adequate levels – domain sensitivity of the Fugl-Meyer Scale (FMS)^{25,27} and Nottingham Sensory Assessment (NSA)¹⁹.

Standardization/test description

NSA identifies sensory deficits through four subscales (Table 1). The total score for tactile sensation (light touch, pressure, pinprick, temperature, tactile location in both hemibodies, simultaneous bilateral tactile touch) ranged from 0 to 90 for the hemibody unaffected, and from 0 to 108 for the affected one; proprioception (execution of movement, direction and joint position) has a maximum of 21 points; stereognosis (recognition of 11 objects) can score from 0 to 22; discrimination between two points (index finger and thenar region) has maximum score of 4 points¹⁹. When it was not possible to test an item, Lima et al.⁹, in the Brazilian version, used a score from 4 to 9, whereas Connell²⁸ gives scores from 4 to 10. The NSA was drafted in 1991, showing good intra-rater reliability and poor inter-rater reliability²⁹. After reduction of items, we found acceptable levels of inter-rater reliability, but they were not good³⁰. The Brazilian version¹⁹ showed high intra- and inter-rater

reliability, except for temperature, and correlation with the sensitivity of the FMS subscale.

The sensitivity of the FMS evaluates the tactile location and sense of movement. From the translation of the manual prepared by Dutil et al.³¹, Michaelsen et al.²⁵ describe tactile sensitivity being evaluated in anterior and posterior region of the shoulder, arm, forearm, thumb, index finger, middle third of the anterior tibial region and plantar region of the foot in both hemibodies, with a maximum of 20 points for upper limb and 4 for lower limb. Sense of movement was evaluated in the shoulder, elbow, wrist, thumb, hip, knee, ankle and hallux, with total score of 8 points for upper limb and 8 for lower limb. In the study by Maki et al.²⁷, the exteroceptive sensitivity was evaluated in the upper limb (without specifying the stimulus location), palm of the hand, thigh and sole of the foot, with a maximum of 8 points. The inter^{25,27} and intra-observer²⁷ sensory domain reliability of the Brazilian version of FMS was excellent.

Table 1. Characteristics of studies and instruments' description

Reference	Characteristics of the Sample	Modality	Material	Forms of Punctuation	Measurement Properties
NSA Lima et al., 2010 ¹⁹	n = 21; Age (years) 49.5±13.6; Time after the stroke (months) 40.2±32.4	Tactile sensation Proprioception Stereognosis Discrimination between two points.	Cotton; test tubes with hot or cold water and talcum powder; monofilament (green); coins of R\$0.01, 0.10 and 1.00; pen; pencil; comb; scissors; sponge; flannel; cup; glass; compass.	Tactile sensation and stereognosis: (0) missing; (1) changed; (2) normal. Discrimination between 2 points: (0) missing; (1) > 3 mm fingers and >8 mm hand; (2) < 3 mm fingers and < 8 mm hand. Proprioception: (0) missing; (1) execution of the wrong motion-direction; (2) direction of motion > 10th; (3) normal or joint position < 10th. (4) to (9) not testable.	<u>Brazilian version</u> ¹⁹ : Inter-rater reliability ICC of 0.80 to 1.00 among the items. Intra-rater reliability ICC of 0.86 to 1.00, except for temperature (ICC - 0.07).
FMS Michaelsen et al., 2011 ²⁵	n=18; Age (years) 59±10; Time after the stroke (months) 39±33.6	Tactile sensitivity and proprioception	Cotton	Tactile sensitivity: (0) lack of sensitivity; (1) hypo- or hypersensitivity (2) normal sensitivity.	<u>Brazilian version</u> : Inter-rater reliability: tactile ICC 0.75; sense of movement ICC of 0.98 ²⁵ Inter-observer reliability: exteroceptive ICC 0.98-0.99; proprioceptive ICC 0.97-0.98.
Maki et al., 2006 ²⁷	n=50; Age (years) 58 (17-81); Time after the stroke (months) 5.(13-127)			Sense of movement (0) does not identify the movement; (1) at least 75% of the correct answers; (2) all the correct answers.	Inter-observer reliability: exteroceptive ICC 0.87 to 0.92; proprioceptive ICC 0.93 to 0.94 ²⁷ .
MTP Faria-Fortini et al., 2011 ²⁴ (Faria, 2008 ²⁶)	n = 55; Age (years) 54±13; Time after the stroke (months) 64±55	Mobile discriminatory touch	Tigre Pincore® brushes references 815/14, 483/18, 183/14 ²⁴ .	Number of correct answers in percentage: 0% indicates that there was no correct answer and 100% indicates that they were all correct.	<u>Original version</u> ¹⁸ Inter and intra-rater reliability: ICC 0.92
Brasil-Neto; Lima, 2008 ²³	n=25; Age (years) 58.2±11.1; Time after the stroke (months) 43.8±55.4		Condor® brushes no. 20 of "light", "medium" and "heavy" "bristles" ²³ .		<u>Brazilian version</u> : Not evaluated

NSA: Nottingham Sensory Assessment; FMS: Fugl-Meyer Scale; MTP: Moving Touch Pressure; ICC: intraclass correlation coefficient

Moving Touch Pressure (MTP) measures the tactile sensitivity, by discriminating the sensation generated by 3 brushes (1 cm wide and 2.5 cm in length) of different textures, and verbal indication of which brush touched the skin. The brush is positioned at an angle of 30° with the surface of the skin, with enough force to slightly bend the bristles, and stimulus applied to two centimeters from proximal to distal in the distal phalanx's palm surface of the index finger in sliding movements^{23,24,26}.

Measurement properties

The studies by Brasil-Neto; Lima, 2008²³ and Faria-Fortini et al., 2011²⁴ were included to satisfy the first goal of this review, i.e. to describe tests available in Portuguese, which are standardized and with quantitative results. However, they did not evaluate the psychometric properties of these tests. Of the included studies, only three assessed measurement properties^{19,25,27} and all used the classical theory. Of the nine items proposed by COSMIN, only reliability was evaluated in three studies. In addition to reliability, the study of Lima et al.¹⁹ also assessed Internal Consistency with Cronbach's alpha and Criterion Validity with Pearson Correlation test.

As for their methodological quality of reliability, of the eleven items evaluated, seven showed excellent quality in all three assessed articles (Table 2). None of the studies had a sample size that reached a number of participants of at least 100 individuals to be considered excellent, according to the criteria of COSMIN. The time interval between the assessments was three days and a week, which is considered inferior than the ideally suggested by COSMIN to achieve an excellent score. Still considering the type of score used by the scales, the Kappa-related items were considered not applicable.

DISCUSSION

The description of the tools' qualities of measurement is critical to improve the quality of the research and to enable reliable data for clinical practice, because the measurement allows the quantitative comparison of the results³². Translation and mensuration studies of measurement properties of sensory evaluation tests are recent in Brazil. Regarding the psychometric property

of reliability, studies of sensory assessment tools showed, generally, good results. Maki et al.²⁷ found excellent inter-observers reliability for the sensitivity of the FMS subscale when an appraiser applied the test, while the other two watched and scored. When the test was applied by two different examiners, Michaelsen et al.²⁵ reported excellent reliability for the tactile and proprioceptive sensitivity after adapting the manual with pictures illustrating the location for the tactile test and the position of the hands for the proprioceptive test. Using the original version of FMS³³, Sanford et al.³⁴ found excellent inter-rater reliability among three physical therapists applying the scale.

Table 2. Evaluation of the reliability's study design according to the COSMIN Checklist (Box B)

Reference	Lima et al., 2010 ¹⁹	Michaelsen et al., 2011 ²⁵	Maki et al., 2006 ²⁷
Test	NSA	FMS	FMS
Items evaluated			
Sample size	P	P	G
Two evaluation measures	E	E	E
Independent evaluation	E	E	E
Interval of described time	E	E	E
Participants' Clinical stability	E	E	E
Interval of described time	R	G	R
Conditions of the application of similar test	E	E	E
There are major flaws in the method's design	E	E	R
Was it calculated the ICC?	E	E	E
Were Kappa calculated for dichotomous/nominal or ordinal items?	NA	NA	NA
Was the weighted Kappa calculated for ordinal scores?	NA	NA	NA

E = excellent; G = good; R = reasonable; P = poor; NA = not applicable

MTP is used in Brazil^{23,24}, but no national study evaluated its psychometric characteristics. It was developed, described and had its measurement properties evaluated by Dannenbaum et al.¹⁸ demonstrating significant correlation with the Semmes-Weinstein and Moberg tests, and high inter- and intra-rater reliability.

The index finger's distal sensitivity assessed through MTP was significant, along with the grip force variables, to explain the levels of activity in the upper limb²⁴.

The Brazilian version of the NSA showed excellent intra-rater reliability (test-retest) and inter-rater in all NSA items.

FMS' sensitivity session is restricted to the assessment of tactile location and sense of movement. NSA, in its turn, along with the tactile sensation and proprioception, also evaluates the stereognosis and the discrimination between two points¹⁹. While in FMS and NSA touch is evaluated statically, in the MTP it is evaluated dynamically, which can be more relevant to the function, however the application is limited in just one location of the hand. FSM offers a quick and less detailed review of the sensitivity, while the NSA offers a detailed one and the MTP offers a specific assessment for manual mobile touch.

A systematic review¹⁷ evaluated seven scales concerning the time of application, cost, need for equipment and portability. In a score from 0 to 10, three scales achieved good score since the modified version of the NSA and sensory subscale of the FSM gained nine points and the MTP eight points.

The Semmes-Weinstein test is used in the sensory evaluation of peripheral nerve injuries³⁵, however the article that evaluated its measurement properties was not found. Although the search in the database has not identified any article with this test, some research already used it in patients affected by stroke^{36,37}.

With the exception of NSA that evaluated internal consistency and criterion validity¹⁹, other studies showed the limitation of assessing just reliability. We highlight the importance of research that evaluate other measurement properties of these instruments. Only the presence of adequate reliability is not enough to prove that the instrument evaluates what it intends to. Sousa³⁸ used the NSA to report improvement in the tactile sensitivity of patients subjected to sensory stimulation of the hand. However, sensitivity to change was not assessed in any of the sensitivity assessment tools available in Portuguese. We suggest that the evaluation of these instruments' ability to detect changes is included, enabling the assessment of the sensory rehabilitation effect in this population.

Considering the measurement properties' lack of assessment of sensory evaluation tools available in

Portuguese – including the properties described by COSMIN –, it was possible to analyze them considering just the methodological quality used to assess reliability, which is a limitation of this study.

CONCLUSION

This review found three sensory evaluation tools in Portuguese that clearly describe the application and present a quantitative score for sensory evaluation of post-stroke patients. Reliability was the only measurement property evaluated in two of the instruments (FMS and NSA) and we considered it adequate. We also evaluated the internal consistency and criterion validity of the NSA. However, future studies need to evaluate the other measurement properties of these instruments. Few sensory tests for the population with stroke are available in Portuguese and with insufficient evaluation of their measurement properties.

REFERENCES

1. Garritano CR, Luz PM, Pires MLE, Barbosa MTS, Batista KM. Análise da tendência da mortalidade por acidente vascular cerebral no Brasil no século XXI. *Arq Bras Cardiol*. 2012;98(6):519-27.
2. Minelli C, Fen LF, Minelli DP. Stroke incidence, prognosis, 30-day, and 1-year case fatality rates in Matao, Brazil: a population-based prospective study. *Stroke*. 2007;38(11):2906-11.
3. Connell LA, Lincoln NB, Radford KA. Somatosensory impairment after stroke: frequency of different deficits and their recovery. *Clin Rehabil*. 2008;22(8):758.
4. Tyson SF, Hanley M, Chillala J, Selley AB, Tallis RC. Sensory loss in hospital-admitted people with stroke: characteristics, associated factors, and relationship with function. *Neurorehabil Neural Repair*. 2008;22(2):166-72.
5. Kim JS, Choi-Kwon S. Discriminative sensory dysfunction after unilateral stroke. *Stroke*. 1996;27(4):677-82.
6. Scalha TB, Miyasaki E, Lima NMFV, Borges G. Correlations between motor and sensory functions in upper limb chronic hemiparetics after stroke. *Arq Neuropsiquiatr*. 2011;69(4):624-9.
7. Carey LM. Somatosensory loss after stroke. *Crit Rev Phys Rehabil Med*. 1995;7(1):51-91.
8. Reding MJ, Potes E. Rehabilitation outcome following initial unilateral hemispheric stroke. *Stroke*. 1988;19(11):1354-8.

9. Han L, Law-gibson D, Reding M. Key neurological impairments influence function-related group outcomes after stroke. *Stroke*. 2002;33:1920-4.
10. Sommerfeld DM, von Arbin MH. The impact of somatosensory function on activity performance and length of hospital stay in geriatric patients with stroke. *Clin Rehabil*. 2004;18(2):149-55.
11. Sullivan JE, Hedman LD. Sensory dysfunction following stroke: incidence, significance, examination, and intervention. *Top Stroke Rehabil*. 2008;15(3):200-17.
12. Wagner JM, Lang CE, Sahrman SA, Hu Q, Bastian AJ, Edwards DF, et al. Relationships between Sensorimotor Impairments and Reaching Deficits in Acute Hemiparesis. *Neurorehabil Neural Repair*. 2006;20(3):406-16.
13. Lin SI. Motor function and joint position sense in relation to gait performance in chronic stroke patients. *Arch Phys Med Rehabil*. 2005;86(2):197-203.
14. Lin PY, Yang YR, Cheng SJ, Wang RY. The relation between ankle impairments and gait velocity and symmetry in people with stroke. *Arch Phys Med Rehabil*. 2006;87(4):562-8.
15. Lee MJ, Kilbreath SL, Refshauge KM. Movement detection at the ankle following stroke is poor. *Aust J Physiother*. 2005;51(1):19-24.
16. Winward CE, Halligan PW, Wade DT. Current practice and clinical relevance of somatosensory assessment after stroke. *Clin Rehabil*. 1999;13(1):48-55.
17. Connell LA, Tyson SF. Measures of sensation in neurological conditions: a systematic review. *Clin Rehabil*. 2011;26(1):68-80.
18. Dannenbaum RM, Michaelsen SM, Desrosiers J, Levin MF. Development and validation of the two new sensory tests of the hand for patients with stroke. *Clin Rehabil*. 2002;16(6):630-9.
19. Lima DHF, Queiroz AP, Salvo G, Yoneyama SM, Oberg TD, Lima NMFV. Versão brasileira da Avaliação Sensorial de Nottingham: validade, concordância e confiabilidade. *Rev Bras Fisioter*. 2010;14(2):166-74.
20. Moher D, Liberati A, Tetzlaff J, Altman DG; PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med*. 2009;151(4):264-9.
21. Terwee CB, Bot SDB, Boer MR, Van Der Windt DAWM, Knol DL, Dekker J, et al. Quality criteria were proposed for measurement properties of health status. *J clin epidemiol*. 2007;60:34-42.
22. Mokkink LB, Terwee CB, Patrick DL, Alonso J, Stratford PW, Knol DJ, et al. COSMIN checklist manual. 2012. Disponível em: www.cosmin.nl
23. Brasil-Neto JP, Lima AC. Sensory Deficits in the unaffected hand of hemiparetic stroke patients. *Cog Behav Neurol*. 2008;21(4):202-5.
24. Faria-Fortini I, Michaelsen SM, Cassiano JG, Teixeira-Salmela LF. Upper Extremity Function in Stroke Subjects: Relationships between the International Classification of Functioning, Disability, and Health Domains. *J Hand Ther*. 2011;24(3):257-65.
25. Michaelsen SM, Rocha AS, Knabben RJ, Rodrigues LP, Fernandes CGC. Translation, adaptation and inter-rater reliability of the administration manual of Fugl-Meyer assessment/Tradução, adaptação e confiabilidade interexaminadores do manual de administração da escala de Fugl-Meyer. *Rev Bras Fisioter*. 2011;15(1):80-8.
26. Faria I. Estudo da função do membro superior em hemiplégicos crônicos [dissertação – mestrado em Ciências da Reabilitação]. Belo Horizonte: Universidade Federal de Minas Gerais. Escola de Educação Física, Fisioterapia e Terapia Ocupacional, 2008.
27. Maki T, Quagliato EMAB, Cacho EWA, Paz LPS, Nascimento NH, Inoue MMEA, et al. Estudo de confiabilidade da Escala de Fugl-Meyer no Brasil. *Rev Bras Fisioter*. 2006;10(2):177-83.
28. Connell LA. Sensory impairment and recovery after stroke [tese]. Nottingham: Universidade de Nottingham, 2007. Disponível em: http://etheses.nottingham.ac.uk/247/1/PHD_final.pdf
29. Lincoln NB, Crow JL, Jackson JM, Waters GR, Adams SA, Hodgson P. The unreliability of sensory assessments. *Clin Rehabil*. 1991;5:273-82.
30. Lincoln NB, Jackson JM, Adams SA. Reliability and revision of the Nottingham sensory assessment for stroke patients. *Physiotherapy*. 1998;84(8):358-65.
31. Dutil E, Arsenault AB, Corriveau H, Prévost R. Protocole d'évaluation de la fonction sensori-motrice: Test de Fugl-Meyer. Montréal: La librairie de l'Université de Montréal; 1989.
32. Gadotti IC, Vieira ER, Magge DJ. Importance and clarification of measurement properties in rehabilitation. *Rev Bras Fisioter*. 2006;10(2):137-46.
33. Fugl-Meyer AR, Jaasko L, Leyman I, Olsson S, Stegling S. The post stroke hemiplegic patient. 1. A method for evaluation of physical performance. *Scand J Rehabil Med*. 1975;7(1):13-31.
34. Sanford J, Moreland J, Swanson LR, Stratford PW, Gowland C. Reliability of the Fugl-Meyer assessment for testing motor performance in patients following stroke. *Phys Ther*. 1993;73:447-54.
35. Bell-Krotowski J, Tomancik E. The repeatability of testing with Semmes-Weinstein monofilaments. *J Hand Surg*. 1987;12:155-61.
36. Silva FSR. Efeito da postura no controle da força de preensão do membro superior não parético ao acidente vascular encefálico [dissertação – pós-graduação *strictu sensu* em Ciências do Movimento]. Florianópolis: Universidade do Estado de Santa Catarina, 2011.
37. Torriani C, Mota EPO, Sales ALM, Ricci M, Nishida P, Marques L, et al. Efeitos da estimulação motora e sensorial no pé de pacientes hemiparéticos pós acidente vascular encefálico. *Rev Neurociênc*. 2008;16(1):25-9.

38. Sousa MJN. Eficácia de um programa de estimulação sensorial na mão em indivíduos com déficits sensoriais, seqüela de AVC [monografia – licenciatura em Fisioterapia].

Barcarena: Escola Superior de Saúde da Universidade Atlântica, 2011.