

SHORT COMMUNICATION

Interaction between ostracods and anurans: a review and new records in Brazil

Matheus de Toledo Moroti,¹ Edélcio Muscat,¹ Mariana Pedrozo,² Iberê Farina Machado,³ Leandro Talione Sabagh,⁴ and Diego José Santana²

¹ Projeto Dacnis. Ubatuba e São Francisco Xavier, SP, Brazil. E-mail: mmoroti@gmail.com.

² Instituto de Biociências, Laboratório de Zoologia, Universidade Federal de Mato Grosso do Sul. Cidade Universitária, 79070-900, Campo Grande, MS, Brazil.

³ Instituto Boitatá. Rua 121, QD F 42-A, LT 12, Nº 126, Loja 1, 74085-480, Setor Sul, Goiânia, GO, Brazil.

⁴ Departamento de Ecologia, Laboratório de Limnologia, Universidade Federal do Rio de Janeiro. Cidade Universitária, 21941-902, Rio de Janeiro, RJ, Brazil.

Keywords: bromeliculous, bromeligenous Crustacea, natural history, phoresy.

Palavras-chave: bromelícola, bromelígena Crustacea, foresia, história natural.

Both ostracods and amphibians are able to colonize a wide variety of freshwater environments (Balian *et al.* 2008, Martens *et al.* 2008, Ottonelo and Romano 2011). However, unlike amphibians, ostracods also are widely distributed in marine environments (Coimbra and Bergue 2011). The first fossil records of ostracods are dated from the Ordovician Period and, in the Carboniferous Period, these crustaceans began to colonize freshwater environments (Coimbra and Bergue 2011). Ostracods are found in all types of aquatic environments, such as lakes, rivers, and humid soils of tropical forests and phytotelmata (Pinto and Purper 1970). Like ostracods, some species of anurans are associated with phytotelmata. For example, some anurans use bromeliads throughout their

life cycle (bromeligenous), whereas others occupy phytotelmata, using them for shelter and foraging places, but do not reproduce in them (bromeliculous) (Peixoto 1995).

The microhabitats formed by the accumulation of water and debris in the bromeliads maintain diverse invertebrate communities (Richardson 1999, Sabagh *et al.* 2012). Among invertebrates, non-marine ostracods typically are found in this kind of environment (Müller 1879, Lopez *et al.* 2002). The success of ostracods in non-marine environments reflects the resilience of the group and its use of other organisms, such as mammals (Lopez *et al.* 2002), birds (Green and Figuerola 2005), reptiles, and amphibians (Lopez *et al.* 2005) as vectors for dispersion.

Phoresy is a type of commensal interaction between two species, in which one organism acts as a vector for the dispersion of another, with no detrimental effects on either organism (Houck and O'Connor 1991). Ecologically, bromeliads

Received 06 August 2018
Accepted 29 April 2019
Distributed December 2019

function as “islands,” representing disconnected freshwater habitats for non-marine ostracods (Frank and Lounibos 1987, Sabagh and Rocha 2014, Lantyer-Silva *et al.* 2016). Given the use of bromeliads by some species of anurans, the frogs are used by ostracods as phoretic vectors (Lopez *et al.* 2005, Sabagh *et al.* 2011). However, such interactions are poorly documented and may not be noticed owing to the small size of these invertebrates. Herein we present five new records of this phoresy between ostracods and four anuran species from four localities in southeastern Brazil, and provide a list of all neotropical amphibians recorded in these relationships.

During fieldwork at localities associated with the Dacnis Project (23.4627° S, 45.1329° W, WGS-84; 37 m a.s.l.) in municipality of Ubatuba and state of São Paulo (Brazil), we found two species of anurans with ostracods on their bodies. The first interaction was recorded on 15 July 2015; the toad *Dendrophryniscus brevipollicatus* Jiménez de la Espada, 1870 (Bufonidae) had an individual of *Elpidium* sp. (Crustacea, Ostracoda) attached to its back, close to its head (Figure 1A). The second interaction was on 09 September 2015 at the same locality. The treefrog *Fritziana mitus* Walker, Wachlevski, Nogueira da Costa, Nogueira-Costa, Garcia, and Haddad, 2018 (Hemiphractidae) had three individuals of *Elpidium* sp. (Crustacea, Ostracoda) attached to its flanks near its legs (Figure 1B). We found both anurans in the central tank of bromeliads.

On 15 August 2015 in the Alcatrazes Archipelago (24.1051° S, 45.6974° W, WGS-84; 134 m a.s.l.), located in the Atlantic Ocean, about 3500 m of the municipality of São Sebastião in state of São Paulo (Brazil), we recorded one individual of *Ololygon alcatraz* (Lutz, 1973) (Hylidae) inside a bromeliad with five ostracods attached to its trunk, foot, and tibia (Figure 1C).

In the shrubs around the “Lago da Mata” area on the campus of Universidade do Vale do Paraíba (UNIVAP) (23.2084° S, 45.9701° W, WGS-84; 591 m a.s.l.) in the municipality of São

José dos Campos and state of São Paulo (Brazil), we found a hylid frog, *Scinax crospedospilus* (Lutz, 1925), with two *Elpidium* sp. attached to its flanks close to its head and legs (Figure 1D) on 21 October 2015. We found the same interaction between these species on 08 January 2018 in the area of the Dacnis Project in São Francisco Xavier (22.8744° S, 45.9306° W, WGS-84; 884 m a.s.l.), a district of the municipality of São José dos Campos in the state of São Paulo (Brazil). Here, we found seven *Elpidium* sp. attached to the flanks near the legs of a *S. crospedospilus*. Both localities were close to a flooded area where we recorded *S. crospedospilus* calling from inside a bromeliad.

All the voucher specimens were deposited in collections (Appendix I). We did not collect the specimen of *O. alcatraz* or the ostracods found on that individual. The species is the only one occurring on the island and is “Critically Endangered” according to IUCN criteria (Rodrigues and Cruz 2004). Ostracods were identified with the descriptions of Pinto and Jocqué (2013) and Pereira (2013). Despite keys to species of *Elpidium*, we only identified them to the genus, because there are records of cryptic diversity and undescribed species (Little and Herbet 1996).

In addition to the new interactions described above, we performed an exhaustive search of the literature for studies that reported interactions between ostracods and amphibians in the neotropics. We also used “Google Scholar” to locate papers published through October 2019 with the combination of the following keywords: “ostracods,” “Amphibia,” “phoresy,” and “interaction.” Amphibian nomenclature is consistent with that of Frost (2019). The publications then were filtered to include only those dealing with neotropical amphibians.

In total, we tallied 20 species of amphibians that interact with ostracods in 13 localities (Table 1). All interactions were observed in Brazil and the ostracods were identified as *Elpidium* sp. The interaction was only recorded for anurans. Hylidae had the largest number of records, with




Figure 1. Interaction between ostracods and anurans of four species in three families. (A) *Dendrophryniscus brevipollicatus* (Bufonidae) had an individual of *Elpidium* sp. (B) The treefrog *Fritziiana mitus* (Hemiphraactidae) with one individual of *Elpidium* sp. (C) The treefrog *Ololygon alcatraz* (Hylidae) had five individuals of *Elpidium* sp. (D) the treefrog *Scinax crospedospilus* (Hylidae) had two individuals of *Elpidium* sp. Dashed circle indicates places with ostracods attached.

16 species, followed by Hemiphraactidae with 2, and Cycloramphidae and Bufonidae with one each.

Prior records of interactions between ostracods and amphibians are mainly from coastal regions (e.g., Lopez *et al.* 2002, 2005, Sabagh and Rocha 2014) between elevations from 0–963 m a.s.l. The two records of ostracods associated with *Scinax crospedospilus* in the Mantiqueira Mountains at two different localities (Table 1) are the the farthest inland at approximately 80 km from the coast.

Our results show that ostracods are mainly associated with arboreal frogs that use

bromeliads. The only species with adherent ostracods that is not arboreal is *Thoropa miliaris* (Spix, 1824) (Sabagh and Rocha 2014). This species usually is found in rocky areas in forests or at the forest edge; the frog deposits its eggs on stones covered by a film of fresh water (Bokermann 1965). *Thoropa miliaris* has been reported to be bromeliculous because the frogs use bromeliads as shelter (Lacerda *et al.* 2009, Pertel *et al.* 2010, Sabagh and Rocha 2014); thus, the species may be a potential disperser of ostracods among aquatic environments.

We observed two populations of *Scinax crospedospilus* calling and sheltering in bromeliads; however, we found neither tadpoles

Table 1. Literature records of phoretic behavior in Brazilian anurans and ostracods and new records of this behavior.

Anuran species	Study area	Altitude (m a.s.l.)	Municipality, Brazilian state	References to the first record
Bufonidae				
<i>Dendrophryniscus brevipollicatus</i>	Projeto Dacnis	37	Ubatuba, São Paulo	This study
Cycloramphidae				
<i>Thoropa miliaris</i>	Parque Estadual Serra da Tiririca	145	Niterói, Rio de Janeiro	Sabagh and Rocha 2014
	MoNa Morro da Urca e Pão de Açúcar	0	Rio de Janeiro, Rio de Janeiro	Sabagh and Rocha 2014
Hemiphractidae				
<i>Fritziana goeldii</i>	Parque Nacional Serra dos Órgãos	963	Teresópolis, Rio de Janeiro	Lopez <i>et al.</i> 2005
<i>Fritziana mitus</i>	Projeto Dacnis	37	Ubatuba, São Paulo	This study
Hylidae				
<i>Aparasphenodon brunoi</i>	Barra de Maricá	6	Rio de Janeiro, Rio de Janeiro	Lopez <i>et al.</i> 2005
	Parque Nacional da Restinga de Jurubatiba	9	Macaé, Rio de Janeiro	Lopez <i>et al.</i> 2005
<i>Aparasphenodon arapapa</i>	Reserva Natural Boa União	95	Ilhéus, Bahia	Lantyer-Silva <i>et al.</i> 2016
<i>Aplastodiscus arildae</i>	Parque Nacional Serra dos Órgãos	963	Teresópolis, Rio de Janeiro	Lopez <i>et al.</i> 2005
<i>Boana albomarginata</i>	Grumari	10	Rio de Janeiro, Rio de Janeiro	Sabagh <i>et al.</i> 2011
<i>Boana semilineata</i>	Parque Nacional da Restinga de Jurubatiba	9	Macaé, Rio de Janeiro	Lopez <i>et al.</i> 2005
<i>Dendropsophus decipiens</i>	Alto da Buchada	200	São Lourenço da Mata, Pernambuco	Araújo <i>et al.</i> 2019
<i>Oloolygon alcatraz</i>	Ilha dos Alcatrazes	134	São Sebastião, São Paulo	This study
<i>Oloolygon littorea</i>	Parque Estadual Serra da Tiririca	145	Niterói, Rio de Janeiro	Sabagh <i>et al.</i> 2011
<i>Oloolygon perpusilla</i>	Barra de Marica	6	Rio de Janeiro, Rio de Janeiro	Lopez <i>et al.</i> 2002
	MoNa Morro da Urca e Pão de Açúcar	0	Rio de Janeiro, Rio de Janeiro	Sabagh <i>et al.</i> 2011

Table 1 - Continued.

Anuran species	Study area	Altitude (m a.s.l.)	Municipality, Brazilian state	References to the first record
<i>Scinax crospedospilus</i>	Lago da Mata (Campus UNIVAP)	591	São José dos Campos, São Paulo	This study
	Projeto Dacnis (SFX)	884	São Francisco Xavier, São Paulo	This study
<i>Scinax auratus</i>	Alto da Buchada	200	São Lourenço da Mata, Pernambuco	Araújo <i>et al.</i> 2019
<i>Scinax cuspidatus</i>	Parque Estadual Serra da Tiririca	145	Niterói, Rio de Janeiro	Sabagh and Rocha 2014
<i>Scinax pachycrus</i>	Alto da Buchada	200	São Lourenço da Mata, Pernambuco	Araújo <i>et al.</i> 2019
<i>Scinax x-signatus</i>	Alto da Buchada	200	São Lourenço da Mata, Pernambuco	Araújo <i>et al.</i> 2019
<i>Sphaenorhynchus aff. surdus</i>	Parque Estadual de Itapeva	33	Torres, Rio Grande do Sul	Colombo <i>et al.</i> 2008
<i>Xenohyla truncata</i>	Barra de Maricá	6	Rio de Janeiro, Rio de Janeiro	Lopez <i>et al.</i> 1999

or eggs at either site. This record suggests that *S. crospedospilus* is not an obligate bromeliad species (bromelicolous), as observed by Sabagh and Rocha (2014) for *Scinax cuspidatus* (Lutz, 1925), as other hylid species (Araújo *et al.* 2019). *Oloolygon alcatraz* reproduces and completes its life cycle in bromeliads (bromeligenous), as do *Fritziana mitus* and *Dendrophryniscus brevipollicatus* (Haddad and Prado 2005, Haddad *et al.* 2013, Menegucci *et al.* 2017). Thus, it is likely that both, bromeligenous and bromelicolous frogs are important vectors for transporting ostracods between bromeliads.

The interaction between amphibians and ostracods is considered phoresy by several authors (e.g., Seidel 1989, Lopez *et al.* 2005, Sabagh *et al.* 2011, Lantyer-Silva *et al.* 2016) or as a prey-predator relationship when tadpoles feed on ostracods (Lopez *et al.* 2002). However, the trophic network in which ostracods and amphibians are involved is complex and a variety

of relationships between these taxa are known today (Ottonelo and Romano 2011). For example, ostracods are naturally found in the digestive tract of tadpoles (Sabagh *et al.* 2012) and can pass unharmed through the amphibian intestine, being eliminated through the feces (Hartmann 1985). Thus, such transport also allows the colonization of new environments (Lopez *et al.* 2002). Conversely, depending on the abundance of these ostracods, they may feed on amphibian eggs and tadpoles (Ottonelo and Romano 2011). Thus, it seems that further studies should be conducted to understand the relationship between amphibians and ostracods, as this interaction apparently is more complex than has been documented in most studies.

In summary, this is the first record of an interaction: (1) between ostracods and an insular anuran (*Oloolygon alcatraz*); (2) involving toads of the family Bufonidae (*Dendrophryniscus brevipollicatus*); and (3) between the treefrog species *Scinax crospedospilus*, *Oloolygon alcatraz*

and *Fritziana mitus*. Last, an updated list of neotropical frogs that interact with ostracods is provided.

Acknowledgments.—The authors thank the Dacnis Project for support. MTM and MP are grateful to CAPES for scholarships. LTS thanks CNPq (#150841/2017-5). We also thank Alejandro Valencia-Zuleta for reading and suggesting on earlier drafts of the manuscript. We thank ICMBio for the licenses granted (SISBIO 48636-2, 48620-2, 64215-2). 

References

- Araújo, A. P., C. M. Bastos, R. V. I. Santos, G. J. B. Moura, M. Melo-Junior and M. S. Tinoco. 2019. Novel records of phoresy among microcrustaceans and bromeliad treefrogs in the Atlantic Rainforest of Northeast Brazil. *Herpetology Notes* 12: 531–535.
- Balian, E. V., H. Segers, C. Lévêque, and K. Martens. 2008. The freshwater animal diversity assessment: an overview of the results. *Hydrobiologia* 595: 627–637.
- Bokermann, W. C. A. 1965. Notas sobre as espécies de *Thoropa* Fitzinger (Amphibia, Leptodactylidae). *Anais da Academia Brasileira de Ciências* 37: 525–537.
- Coimbra, J. C. and C. T. Bergue. 2011. Ostracodes. Pp 35–49 in I. S. Carvalho (ed.), *Paleontologia: Microfósseis, Paleoinvertebrados*. Vol. 2. 3th Edition. Rio de Janeiro Editora Interciência.
- Colombo, P., A. Kindel, G. Vinciprova, and L. Krause. 2008. Composição e ameaças à conservação dos anfíbios anuros do Parque Estadual de Itapeva, município de Torres, Rio Grande do Sul, Brasil. *Biota Neotropica* 8: 229–240.
- Frank, J. H. and L. P. Lounibos. 1987. Phytotelmata: swamps or islands? *Florida Entomologist* 70: 14–20.
- Frost, D. R. (ed.). 2019. Amphibian Species of the World: an Online Reference. Version 6.0. Electronic Database accessible at <http://research.amnh.org/herpetology/amphibia/index.html>. American Museum of Natural History, New York, USA. Captured on 3 October 2019.
- Green, A. J. and J. Figuerola. 2005. Recent advances in the study of long-distance dispersal of aquatic invertebrates via birds. *Diversity and Distribution* 11: 149–156.
- Haddad, C. B. F. and C. P. A. Prado. 2005. Reproductive modes in frogs and their unexpected diversity in the Atlantic Forest of Brazil. *BioScience* 55: 207–217.
- Haddad, C. F. B., L. F. Toledo, C. P. A. Prado, D. Loebmann, J. L. Gasparini and I. Sazima. 2013. *Guia dos Anfíbios da Mata Atlântica: Diversidade e Biologia*. São Paulo. Anolis Books Editora. 544 pp.
- Hartmann, G. 1985. Ostracoden aus der Tiefsee des Indischen Ozeans und der Iberischen See sowie von ostatlantischen sublitoralen Plateaus und Kuppen. Mit einer Tabelle der bislang bekannten rezenten Tiefseeostracoden. *Senckenb Marit* 17: 89–146.
- Houck, M. A. and B. M. O'Connor. 1991. Ecological and evolutionary significance of phoresy in the Astigmata. *Annual Review of Entomology* 36: 611–636.
- Lacerda, J. V. A., B. Assis, D. J. Santana, and R. N. Feio. 2009. Anurans in bromeliads, Parque Estadual da Serra do Brigadeiro, state of Minas Gerais, southeastern Brazil. *Check List* 5: 800–806.
- Lantyer-Silva, A. S. F., C. C. Souza, I. Jaloretto, and M. Solé. 2016. *Aparasphenodon arapapa* (Bahia's Broad-snout Casque-headed Treefrog). Phoretic Ostracods. *Herpetological Review* 47: 106.
- Little, T. J. and P. D. N. Hebert. 1996. Endemism and ecological islands: the ostracods from Jamaican bromeliads. *Freshwater Biology* 36: 327–338.
- Lopez, L. C. S., P. P. Rodrigues, and R. I. Rios. 1999. Frogs and snakes as phoretic dispersal agents of bromeliad ostracods (Limnocytheridae: *Elpidium*) and annelids (Naididae: *Dero*). *Biotropica* 31: 705–708.
- Lopez, L. C. S., B. Filizona, I. Deiss and R. I. Rios. 2005. Phoretic behaviour of bromeliad annelids (*Dero*) and ostracods (*Elpidium*) using frogs and lizards as dispersal vectors. *Hydrobiologia* 549: 15–22.
- Lopez, L. C. S., D. A. Gonçalves, A. Mantovani, and R. I. Rios. 2002. Bromeliad ostracods pass through amphibian (*Scinax perpusillus*) and mammalian guts alive. *Hydrobiologia* 485: 209–211.
- Martens, K., I. Schön, C. Meisch, and D. J. Horne. 2008. Global diversity of ostracods (Ostracoda, Crustacea) in freshwater. *Hydrobiologia* 595: 185–193.
- Menegucci, R. C., R. Gaiga, and I. F. Machado. 2017. *Fritziana ulei* (Miranda-Ribeiro, 1926): geographic extension, with comments on the natural history of this species. *Check List* 13: 2061.
- Müller, F. 1879. Descrição do *Elpidium bromelium* crustáceo da família dos Cytherídeos. *Arquivos do Museu Nacional do Rio de Janeiro* 4: 27–34.
- Otonello, D. and A. Romano. 2011. Ostracoda and Amphibia in temporary ponds: who is the prey? Unexpected trophic relation in a mediterranean freshwater habitat. *Aquatic Ecology* 45: 55–62.

- Peixoto, O. L. 1995. Associação de anuros a bromeliáceas na Mata Atlântica. *Revista da Universidade Rural, Série Ciências da Vida* 17: 75–83.
- Pereira, E. V. 2013. Taxonomia e ocorrência das espécies do gênero bromelícola *Elpidium* (Crustacea: Ostracoda) em áreas de Mata Atlântica do estado de Santa Catarina, Brasil. Unpublished M.Sc. Dissertation. Instituto de Biociências da Universidade de São Paulo, Brazil. 204 pp.
- Pinto, I. D. and I. Purper. 1970. A Neotype for *Elpidium bromeliarum* Müller, 1880:(type species for the Genus) and a revision of the Genus *Elpidium* (Ostracoda). Unpublished M.Sc. Dissertation. Universidade Federal do Rio Grande do Sul, Brazil.
- Pinto, R. L. and M. Jocqué. 2013. A new species of *Elpidium* (Crustacea, Ostracoda) from bromeliads in Cusuco National Park, Honduras. *ZooKeys* 313: 45–59.
- Pertel, W., R. L. Teixeira, and R. B. Ferreira. 2010. Comparison of diet and use of bromeliads between a bromelicolous and a bromeligenous anuran at an Inselberg in the southeastern of Brazil. *Caldasia* 32: 149–159.
- Richardson, B. A. 1999. The bromeliad microcosm and the assessment of faunal diversity in a neotropical forest. *Biotropica* 31: 321–336.
- Rodrigues M. T. and C. A. G. Cruz. 2004. *Scinax alcatraz*. The IUCN Red List of Threatened Species 2004. Electronic Database accessible at <http://dx.doi.org/10.2305/IUCN.UK.2004.RLTS.T55924A11393985.en>. Captured on 19 February 2019.
- Sabagh, L. T. and C. F. D. Rocha. 2014. Bromeliad treefrogs as phoretic hosts of ostracods. *Naturwissenschaften* 101: 493–497.
- Sabagh, L. T., G. L. Ferreira, C. W. C. Branco, C. F. D. Rocha, and N. Y. N. Dias. 2012. Larval diet in bromeliad pools: a case study of tadpoles of two species in the genus *Scinax* (Hylidae). *Copeia* 2012: 683–689.
- Sabagh, L. T., R. J. P. Dias, C. W. C. Branco, and C. F. D. Rocha. 2011. News records of phoresy and hyperphoresy among treefrogs, ostracods, and ciliates in bromeliad of Atlantic forest. *Biodiversity and Conservation* 20: 1837–1841.
- Seidel, B. 1989. Phoresis of *Cyclocypris ovum* (Ostracoda, Podocopida, Cypridae) on *Bombina variegata variegata* (Anura) and *Triturus vulgaris* (Urodela). *Crustaceana* 57: 171–176.

Editor: Jaime Bertoluci

Appendix I. Vouchers for anuran species collected.

Individuals were deposited in the following collections: Museu de Zoologia “Coleção Zoológica da UFG”, Universidade Federal de Goiás, Goiânia, Goiás state, Brazil (*Dendrophyniscus brevipollicatus*: ZUFG-AMP 14277); Coleção Zoológica de Referência, Universidade Federal de Mato Grosso do Sul, Campo Grande, Mato Grosso do Sul state, Brazil (*Scinax crosopedopilus*: ZUFMS-AMP 12522–12523); Coleção Célio Fernando Baptista Haddad, Universidade Estadual de Rio Claro, Rio Claro, São Paulo state, Brazil (*Fritziana mitus*: CCFBH-AMP 40020).