

doi: <http://dx.doi.org/10.11606/issn.1679-9836.v97i2p594-595>

LETTER TO THE EDITOR

Bone microarchitecture evaluation in ankylosing spondylitis patients assessed by lumbar spine trabecular bone score and HR-pQCT

**Pedro Luis Furlam, Valéria de Falco Caparbo,
Carla Gonçalves Schain Saad, Rosa Maria Rodrigues Pereira**

Our article entitled “*Bone Impairment Assessed by Lumbar Spine Trabecular Bone Score (TBS) and HR-pQCT in Male Ankylosing Spondylitis Patients*” has the purpose to study bone parameters of ankylosing spondylitis patients assessed by different structural variables than the traditional Dual-Energy X-Ray Absorptiometry.

Ankylosing spondylitis (AS) is a disease that affects the cortical and the trabecular bone compartment in distinct ways, syndesmophytes formation and ligamentous calcification leads to a false idea of bone quality¹, overestimating areal bone mineral density (aBMD) DXA and hiding bone damage caused by the systemic inflammation^{2,3}.

Due to this limitation of using aBMD DXA in AS patients, new techniques, as Trabecular Bone Score (TBS)⁴ and HR-pQCT^{5,6} show up as important alternatives to better assess bone quality in this patients, preventing them from having bone fractures.

TBS is a parameter measured during the lumbar

spine DXA scanning (iNsight Software) that excludes syndesmophytes and other ligamentous calcifications, giving a more reliable picture of bone impairment⁷. HR-pQCT, in other hand, is a High Resolution peripheral Tomography that shows microarchitecture and biomechanical parameters of bone, analyzing distal radius and tibia^{8,9}.

In our study, we evaluated ankylosing spondylitis patients comparing them to age and gender-matched healthy controls. We recorded individual characteristics as age, physical activity, Body Mass Index (BMI), smoking, alcohol intake and history of fracture, statistically adjusting any parameters that showed difference between the compared groups.

By analyzing the data, we concluded that TBS and HR-pQCT are superior methods to differentiate between AS patients and healthy controls compare to aBMD DXA and therefore should be used to evaluate bone quality in AS patients.

REFERENCES

1. Sieper J, Braun J, Rudwaleit M, Boonen A, Zink A. Ankylosing spondylitis: an overview. *Ann Rheum Dis*. 2002;61(Suppl 3):iii8-18.
2. Lee YS, Schlotzhauer T, Ott SM, van Vollenhoven RF, Hunter J, Shapiro J, et al. Skeletal status of men with early and late ankylosing spondylitis. *Am J Med*. 1997;103(3):233-4.
3. Magrey MN, Lewis S, Asim Khan M. Utility of DXA scanning and risk factors for osteoporosis in ankylosing spondylitis-a prospective study. *Semin Arthritis Rheum*. 2016;46(1):88-94.
4. Kolta S, Briot K, Fechtenbaum J, Paternotte S, Armbrecht G, Felsenberg D, et al. TBS result is not affected by lumbar spine osteoarthritis. *Osteoporos Int*. 2014;25(6):1759-64.
5. Klingberg E, Lorentzon M, Göthlin J, Mellström D, Geijer M, Ohlsson C et al. Bone microarchitecture in ankylosing spondylitis and the association with bone mineral density, fractures, and syndesmophytes. *Arthritis Res Ther*. 2013;15(6):R179.
6. Geusens P, Chapurlat R, Schett G, Ghasem-Zadeh A, Seeman E, de Jong J, et al. High-resolution in vivo imaging of bone and joints: a window to microarchitecture. *Nat Rev Rheumatol*. 2014;10(5):304-13.
7. Muschitz C, Kocijan R, Haschka J, Pahr D, Kaider A, Pietschmann P, et al. TBS reflects trabecular microarchitecture in premenopausal women and men with idiopathic osteoporosis and low-traumatic fractures. *Bone*. 2015;79:259-66.
8. Alvarenga JC, Fuller H, Pasoto SG, Pereira RM. Age-related reference curves of volumetric bone density, structure, and biomechanical parameters adjusted for weight and height in a population of healthy women: an HR-pQCT study. *Osteoporos Int*. 2017;28(4):1335-46.
9. Boutroy S, Bouxsein ML, Munoz F, Delmas PD. In vivo assessment of trabecular bone microarchitecture by high resolution peripheral quantitative computed tomography. *J Clin Endocrinol Metab*. 2005;90:6508-15.

Received: October 8, 2018.

Accepted: October 09, 2018.