

DEPARTAMENTO DE INDÚSTRIA, INSPEÇÃO E CONSERVAÇÃO DE PRODUTOS  
ALIMENTÍCIOS DE ORIGEM ANIMAL

Director: Prof. Paschoal Mucciolo

DIVISION OF BACTERIOLOGY, AMERICAN MEAT INSTITUTE FOUNDATION,  
UNIVERSITY OF CHICAGO, ILLINOIS

Director: Dr. C. F. Niven, Jr.

## EFFECT OF BIOTIN ANTAGONISTS AND RELATED SUBSTANCES UPON THE GROWTH OF THE COAGULASE-NEGATIVE STAPHYLOCOCCI (\*)

P. MUCCILO (\*\*)

C. F. NIVEN, JR.

One of the most important public health problems confronting the food industry today is bacterial food poisoning caused by certain members of the genus *Staphylococcus*. Under favorable conditions these organisms may elaborate in a food a toxic substance known as "enterotoxin". Foods containing sufficient quantities of enterotoxin when consumed may produce the commonly encountered symptoms of acute prostration, diarrhea, and vomiting. Although rarely fatal, this type of food poisoning attracts a great deal of public attention, especially when large numbers of persons are affected at one occasion.

The laboratory investigations of food poisoning outbreaks have a number of limitations, and a great deal of the information concerning of the responsible food item in a particular food poisoning outbreak is dependent upon epidemiological studies rather than laboratory examination of the foods consumed. The reasons for these limitations are many, including the fact that no adequate laboratory test exists that will differentiate an enterotoxigenic *Staphylococcus* strain from one that produces no toxin.

In recent years work by EVANS (1948), EVANS and NIVEN (1950), and EVANS, BUETTNER and NIVEN (1950) has definitely indicated that all enterotoxigenic *Staphylococcus* strains are confined to one homogeneous group, namely, the coagulase-positive strains. Not all coagulase-positive staphylococci are capable of producing the toxin, as detected by rhesus monkey feeding tests, but from

---

(\*) Presented at the First Pan American Congress of Veterinary Medicine, Peru — Lima, October 20 to 28, 1951.

(\*\*) Recipient of a Rockefeller Foundation fellowship.

the results of EVANS et al., it would appear that approximately 75 per cent of such strains are enterotoxigenic according to the techniques employed.

It is known that a great deal of growth of a food-poisoning *Staphylococcus*, thus resulting in a very large population, must take place in a food item before sufficient enterotoxin is produced to cause illness. It then appears that a helpful laboratory test to be performed upon suspected foods would be the quantitative determination of the numbers of coagulase-positive staphylococci present in the food. At the moment, however, no such laboratory method is available to obtain this information. At best, one can only determine by very crude methods the approximate numbers of such microorganisms in the sample examined. The reason for this dilemma is the fact that other saprophytic staphylococci are invariably encountered in foods that have physiological characteristics similar to the coagulase-positive group, thus preventing a differentiation directly on the plating medium, and making it necessary to isolate individual colonies for positive identification.

In attempts to develop differential or selective plating media for the quantitative detection of coagulase-positive staphylococci, many techniques have been employed, most of which have relied mainly upon some outstanding characteristic that is possessed by this group of bacteria, such as mannitol or lactose fermentation, blood hemolysis, gelatin liquefaction, or tolerance to various chemicals or unfavorable environmental conditions. Since these attempts have failed, other peculiarities of the coagulase-positive staphylococci have been sought.

Recently, GRETLE (1950), and GRETLE and EVANS (1951) have reported a remarkable difference existing between the coagulase-positive staphylococci and the closely related saprophytes with respect to their biotin requirements. Of the 90 coagulase-positive strains tested none required biotin for growth in a simplified casein hydrolysate medium. On the other hand, of the 46 coagulase-negative strains tested 38 required biotin for growth while the remaining 8 strains were greatly stimulated by this vitamin. Virtually all of these coagulase-negative strains had been isolated from foods, some of which had been strongly implicated by circumstantial evidence as the cause of food poisoning outbreaks. By animal feeding tests none of these coagulase-negative staphylococci appeared to be able to produce enterotoxin.

It would then seem possible to devise a plating, or liquid medium, containing an anti-biotin substance in which the growth of the coagulase-negative staphylococci would be inhibited, but yet would have no effect upon the growth of the coagulase-positive group. This medium would then allow a clear-cut method of quantitatively enumerating the coagulase-positive staphylococci without inter-

ference from the saprophytic strains. This investigation deals with the biotin nutrition of the coagulase-negative staphylococci with the above-stated aim in mind.

#### EXPERIMENTAL

*Cultures:* All of the cultures used in this study were obtained from the collection of the Division of Bacteriology, American Meat Institute Foundation. These cultures had been isolated from foods suspected of having caused food poisoning outbreaks, but no evidence of their enterotoxigenicity has been obtained by animal feeding tests. The physiological characteristics of most of these cultures have been reported by EVANS and NIVEN (1950).

*Techniques:* Throughout most of the work the cultures were carried by serial transfer in a trypticase soy broth (BBL) supplemented with 1 gram of sodium citrate and 0.2 grams of cystine per liter. In order to prepare an inoculum for the simplified media used in the experiments, a 24-hour culture of the organism was centrifuged, washed once with water and then diluted 1:100 in sterile distilled water. One drop of this diluted suspension was used as an inoculum for each of the tubes in the various experiments.

The biotin-free basal medium used throughout this study is listed in Table 1. This medium was identical with that employed by GRETHER (1950) in his studies on the nutrition of these staphylococci. For each of the experiments the medium was prepared from stock solutions of the various constituents that were preserved with 1 per cent of a volatile preservative solution described by HUTNER and BJERKNES (1948). All of the stock solutions were stored in the refrigerator.

All glassware used in these studies was thoroughly washed, soaked in a dichromate cleaning solution overnight, thoroughly rinsed with tap and deionized water, and then dried. The tubes of medium were capped with aluminum caps especially designed for that purpose. These caps were routinely cleaned by boiling in deionized water, followed by a thorough rinse in deionized water.

Most of the studies were conducted with the use of 18 mm. pyrex test tubes especially calibrated as to diameter, and candled to eliminate serious errors with respect to optical density.

Double strength basal medium was added to the tubes in 5 ml. quantities and various increments of the test substances added along with distilled water to bring the final volume to 10 ml. Before inoculating, the tubes were autoclaved for 15 minutes at 121°C.

The tubes were incubated at 30°C. and growth was quantitatively determined at the specified interval of time. Growth was estimated turbidimetrically with aid of a Coleman Spectrophotometer, model II, set at 660 m $\mu$ . and equipped with a PC-5 filter. An uninoculated tube of medium was used as a reference to give an optical density of zero. Readings were recorded in terms of optical density multiplied by 100 in order to dismiss with decimals.

### RESULTS

The substances tested throughout this study were avidin, homobiotin, desthiobiotin and biocytin. Results obtained with each of these substances will be discussed in turn.

*Avidin:* — This substance obtained from egg white is said to be a basic protein which combines stoichiometrically with biotin, thus depriving the test microorganism or the animal of this essential vitamin. Thus, in a basal medium having a constant concentration of biotin, increasing increments of avidin would be expected to inversely affect growth of the test organism resulting a straight line effect. This effect has been demonstrated with many biotin-requiring microorganisms.

On the other hand, GRETLE (1950) reported that a stoichiometric relationship did not exist between avidin and biotin when tested with a biotin-requiring, coagulase-negative *Staphylococcus*, strain H17C. Growth of this microorganism appeared to be constant up to a critical level of avidin, beyond which growth abruptly ceased. If still higher concentrations of avidin were added to the medium, growth again resulted; the amount directly depended upon the concentration of avidin present.

In the present study attempts were made to confirm GRETLE's observations. Results of a typical experiment are given in Table 2. The basal medium contained just sufficient biotin (2 m $\mu$ g./10 ml.) to allow maximum growth. Since pure avidin was not available, a bacteriologically filtered egg white preparation was used as a crude source of this substance. Dilutions of the sterile raw egg white were added aseptically to the previously sterilized basal medium.

As shown in Table 2, the phenomenon of GRETLE was confirmed. Growth of the test organism was unaffected up to a level of 0.15 per cent. Just beyond this concentration growth of the test organism was greatly reduced. However, if still higher levels of egg white were added, growth again increased up to approximately that occurring in the control tube.

In the presence of 20 millimicrograms of biotin per 10 ml. of medium egg white did not affect growth in any concentration tested. The reason for the lack of a stoichiometric relationship between avidin and biotin among the coagulase-negative staphylococci is unknown. Obviously, it would be desirable to repeat these experiments using purified avidin preparations. Growth in the presence of very high levels of egg white as shown in Table 2 is believed to be due to the presence of "biotin-sparing" substances in the egg white that are unaffected by avidin.

Although of great interest, these results would indicate that avidin would have little value in the development of a plating medium for the quantitative detection of coagulase-positive staphylococci.

*Desthiobiotin:* — Structurally, desthiobiotin is similar to biotin with the exception of the removal of the sulfur atom, thus opening the cyclic structure of that part of the molecule. It has been variously reported as a biotin active substance for some test microorganism, as well as having anti-biotin properties for other test organisms (STOCKES and GUNNESS, 1945; RUBIN, DREKTER and MAYER, 1945).

In the present experiments desthiobiotin was found to replace biotin for 17 of the 18 coagulase-negative staphylococci tested. The results are recorded in Table 3. For culture S19 desthiobiotin appeared to have weak anti-biotin properties. No growth occurred during the first three days' incubation in those tubes containing this substance but further incubation resulted in growth in all tubes. Although the exact molar inhibitory ratio between desthiobiotin and biotin was not determined for this microorganism, it would be expected to be very high. As shown in Table 3, 2 millimicrograms of biotin completely reversed the growth inhibition of 200 micrograms of desthiobiotin.

Subsequent experiments indicated that the concentration of desthiobiotin added to the medium in the above-cited experiments was far higher than was necessary. For example, in comparison to 0.04 millimicrograms of biotin per 10 ml. being necessary for half-maximum growth for culture H3A, it was found that 0.1 millimicrogram of desthiobiotin would afford the same amount of growth. A similar concentration of desthiobiotin was found to be necessary for half-maximum growth for culture H2C. These quantitative estimations were conducted under aerobic conditions (10 ml. of medium in 125 ml. erlenmeyer flasks) since quantitative growth response was difficult to achieve for many of the microorganisms in the deep culture test tubes ordinarily employed. The experiments

with desthiobiotin were further complicated by the apparent spontaneous decomposition of this substance when held in aqueous solutions.

*Homobiotin*: — This substance is a chemical homolog of biotin containing 6 carbons in the aliphatic side chain rather than the 5 carbons found in the biotin molecule. Homobiotin has been reported by GOLDBERG et al. (1947), and RUBIN and SCHEINER (1949), to have very potent anti-biotin properties for certain yeast and *Lactobacillus* strains. On the other hand, BELCHER and LICHSTEIN (1949) reported that this substance may replace biotin for other yeast strains.

In the present study 17 of the 18 *Staphylococcus* strains studied were able to use homobiotin as a biotin source. One strain (S36) was found to be strongly inhibited by this substance. In contrast to desthiobiotin for strain S19, 2 millimicrograms of biotin failed to counteract the anti-biotin properties of 200 micrograms of homobiotin for this culture, thus indicating a low molar inhibition ratio between the two substances. (The molar inhibition ratio is defined as the moles of antibiotin required to inhibit one mole of biotin.)

Table 4 lists the growth response obtained with strain H17C when increasing increments of homobiotin were added to the basal medium. As was usually the case, a moderate amount of growth of this strain usually occurred in the basal medium alone, thus indicating a small contamination of biotin in the medium or a carryover of this growth factor in the inoculum. However, small quantities of homobiotin completely inhibited growth. On the other hand, larger quantities of homobiotin afforded growth that equalled maximum growth obtained with biotin. No explanation can be offered for this apparent antibiotin property of homobiotin in small concentrations in contrast to its growth stimulation in higher concentrations. As shown in Table 4, the inhibitory effect of this substance could not be demonstrated in any concentration tested in the presence of 2 millimicrograms of biotin per 10 ml. of medium. The experiment was repeated with only 0.5 millimicrogram of biotin added per 10 ml. of medium. Even at this concentration no antibiotin effect of homobiotin could be detected in any of the levels tested.

The question arises as to whether homobiotin is being used as such by these microorganisms or is first being converted into biotin. BELCHER and LICHSTEIN (1949) obtained rather definite evidence that the homolog was not being converted into biotin before being used by certain yeast strains.

In an attempt to throw some light on this question the relative activities of homobiotin for 8 different strains of staphylococci were tested. As shown

in Table 5, the quantity of this homolog required for half-maximum growth was relatively constant among the individual strains. Other than cultures H3A and H2C (these were run in erlenmeyer flasks) the determinations were made in the 18 mm. test tube cultures as previously described. Under these conditions precise estimation for half-maximum growth could not be made and it is believed that the variations cited, other than culture H10A, could be laid to experimental error. Culture H10A also required unusually high amounts of biotin, thus indicating that no significance could be attached to its high homobiotin requirements. This culture will be discussed more thoroughly in a different section.

These results are rather difficult to interpret. If homobiotin were first converted to biotin before being utilized, the rates of conversion might be expected to differ widely among different strains and thus be reflected in different activities among the cultures. Among the cultures tested this was not the case. On the other hand, a constancy in activity of the homolog as was apparently obtained might merely indicate a small contamination of biotin in the homobiotin, itself. Cultures H3A and D2A required .03-.05 millimicrograms of biotin per 10 ml. for half-maximum growth, thus indicating that biotin, itself, was 1000-1500 times as active as homobiotin. A contamination of the homolog with less than 0.1 per cent biotin could then explain the results obtained. On the other hand, an antibiotin effect of homobiotin, as illustrated in Table 4, could not be demonstrated if it were contaminated with biotin. It is therefore tentatively concluded that these organisms are using homobiotin as such in their metabolism without converting it into biotin.

*Biocytin*: — WRIGHT et al. (1950a) reported the isolation in crystalline form of a component of yeast extract, termed biocytin, that was available as a source of biotin to *Lactobacillus casei* but not to *Lactobacillus arabinosus*. Acid or alkaline hydrolysis yielded free biotin. Further characterization of this substance (WRIGHT et al. [1950b]) revealed that it served as a source of biotin for a number of different microorganisms; that it was stable to a number of commercial enzyme preparations; that it was combinable with avidin, and, that it was relatively inactive as compared to biotin in reversing the bacteriostatic activity of certain biotin antimetabolites. The purified crystalline substance contains about 65 per cent biotin.

Although biocytin has not been described as having antibiotin properties for any microorganisms tested, it was thought desirable to test the biotin activity

of this substance upon the staphylococci used in this study. As shown in Table 3, biocytin in the concentration of 4 millimicrograms per 10 ml. effectively replaced the biotin requirements for the 18 strains tested. Table 6 demonstrates that on a weight basis biotin is from 1.4 to 2.0 times as active as biocytin for the three strains tested. It would, therefore, seem, that within the limits of error biotin and biocytin have equal activities for these microorganisms when calculated on a molar basis.

Mention has been made that culture H10A acquired unusually high concentrations of biotin for growth. This is illustrated in Table 3 wherein it is shown that neither 2  $\mu\text{g}$ . of biotin, or 4  $\mu\text{g}$ . of biocytin per 10 ml. of medium were sufficient to influence growth. Of additional interest is the fact that the culture appears to have limited abilities to synthesize biotin. In repeated experiments, including serial subculture, this organism always grew to a slight extent in the basal medium. This small amount of growth could not have resulted from biotin contamination in the basal medium. This is amply demonstrated by the fact that this organism required at least a 100-fold quantity of biotin for half-maximum growth as did cultures H3A and D2A. These results are presented in Table 6. Accordingly, a similar increase in biocytin concentration was necessary for half-maximum growth. Table 5 also illustrates the high requirement for homobiotin for this organism.

These results would indicate that this culture has an unusually high demand for biotin in its metabolism. Further study of this culture may prove to be fruitful.

*Oleic acid:* — During the past few years it has become generally accepted that a combination of oleic acid and aspartic acid will substitute for biotin for those organisms that require this vitamin. However, oleic acid is, itself, very toxic for most microorganisms, and the range of concentration that can be added to a medium without encountering growth inhibition is very narrow. In order to avoid the toxic effects of oleic acid, it is usually added to the medium in the form of Tween 80. This substance (polyoxyethylene sorbitan mono-oleate) satisfies the oleic acid requirements of various microorganisms tested but does not display in toxicity.



Although aside from the general problem at hand, an attempt was made to determine whether the coagulase-negative staphylococci would also grow in a biotin free medium when supplied with Tween 80. As shown in Table 3, all but one strain grew satisfactorily in such a medium. Strain S36 was unable to grow in this medium. Even after incubation for six days growth in the medium was no better than that occurring in the control. A further study concerning the relationship of oleic acid and biotin in the nutrition of this culture would seem to be warranted. Lack of time did not permit this study.

#### DISCUSSION

No attempt was made to incorporate any of the above-mentioned substances into a selective or differential plating medium for the purpose of detecting the coagulase-positive staphylococci in foods and other materials. These studies would indicate that each of the antibiotin substances tested would be of little or no value in inhibiting the coagulase-negative strains in such a medium. It should be realized, however, that the substances having anti-biotin properties were far from exhausted in this study, and that one may yet be found that will be of value in achieving the goal sought.

In spite of the failure to achieve the practical results desired, it would seem that this study has divulged a number of peculiarities in the biotin metabolism of a few of the *Staphylococcus* strains. Further study of these apparent anomalies might be of value in contributing to our general knowledge of bacterial physiology.

#### SUMMARY

The biotin activity, or anti-biotin properties of a number of substances, was tested on a group of biotin-requiring, coagulase-negative *Staphylococcus* strains. Of the 18 cultures tested desthiobiotin served as a source of biotin for 17 strains. This substance appeared to be 40 per cent as active as biotin in supporting growth for two of the cultures tested. This substance possessed antibiotin properties for one strain (S19).

Homobiotin, one of the chemical homologs of biotin, served as a biotin source for 17 of the 18 strains. Its activity appeared to be less than 0.1 per

cent of that of biotin among the 8 strains tested. Although it replaced biotin in high concentrations, small quantities appeared to possess anti-biotin properties for one of the *Staphylococcus* strains. Homobiotin acted as an anti-biotin substance for strain S36 in all concentrations tested.

In contrast to studies reported on other biotin-requiring microorganisms, there appeared to be no stoichiometric relationship between biotin and avidin in the nutrition of *Staphylococcus*, strain H17C. Upon adding increasing increments of egg white to a medium containing a limited amount of biotin, growth was unaffected until a critical concentration of avidin was reached. At this level little or no growth of the microorganism occurred. At higher levels growth again occurred, the amount depending directly upon the level of egg white added.

All *Staphylococcus* cultures tested (18 strains) were able to utilize biocytin in place of biotin. On a molar basis this substance appeared to have an activity equal to that of biotin.

Strain S36 failed to grow in a biotin deficient medium to which 0.1 per cent Tween 80 was added. All other strains tested were able to grow under these conditions.

In comparison with other strains culture H10A appeared to require approximately 100 times as much biotin for growth. This high requirement could be demonstrated in spite of the fact that the culture was able to grow to a limited degree in the absence of added biotin. This culture also demonstrated similar high requirements for biocytin, homobiotin, and desthiobiotin in a biotin free medium.

#### ACKNOWLEDGEMENTS

The authors wish to thank Dr. S. H. RUBIN, of Hoffmann-La Roche, Inc., Nutley, New Jersey, for the homobiotin used in this study. Appreciation is also expressed to Dr. L. D. WRIGHT, of Sharpe & Dohme, Inc., Glenolden, Pennsylvania, for the generous sample of biocytin.

TABLE 1  
Basal Medium

	Mg per 100 ml
Acid hydrolyzed casein (GBI) .....	500
DL-Tryptophan .....	10
L-Cystine .....	10
Glucose .....	1000
K <sub>2</sub> HPO <sub>4</sub> .....	500
Sodium citrate .....	500
MgSO <sub>4</sub> · 7H <sub>2</sub> O .....	80
NaCl .....	4
FeSO <sub>4</sub> · 4H <sub>2</sub> O .....	4
MnCl <sub>2</sub> · 4H <sub>2</sub> O .....	15
Adenine SO <sub>4</sub> .....	1
Guanine HCl .....	1
Uracil .....	1
Xanthine .....	1
	μg per 100 ml
Thiamine HCl .....	100
Nicotinic acid .....	500
Riboflavin .....	100
Pyridoxine HCl .....	100
Ca pantothenate .....	100
Folic acid .....	1

TABLE 2  
Effect of Different Concentrations of Raw Egg White upon the Growth of *Staphylococcus* sp. H17C in a Medium Containing 2 mμg Biotin per 10 ml.

Raw egg white (per cent)	Growth, 47 hrs. (optical density x 100)
0	130
.02	110
.04	110
.06	110
.08	110
.10	110
.15	36
.20	23
.25	37
.30	52
.40	51
.50	62
1.0	62
5.0	110

TABLE 3

Growth Response of Coagulase-Negative *Staphylococcus* Strains Toward Biotin and Related Substances. Readings are Recorded in Terms of Optical Density x 100.

	Basal	Biotin .002 μg./10ml.	Desthiobiotin 200 μg./10ml.	Desthiobiotin + Biotin	Homobiotin 200 μg./10ml.	Homobiotin + Biotin	Bioctyn .004 μg./10ml.	Tween 80 0.1 per cent
H3A	4	24	—	—	—	—	39	48
H12A	19	48	68	56	60	62	55	54
D2A	1	42	34	45	48	45	48	28
H9A	23	51	62	71	88	53	53	43
H8A	43	88	90	94	92	88	98	90
20-10	23	90	58	63	75	95	75	120
H13A	7	37	33	42	45	37	35	55
S19	27	75	0	83	86	79	72	65
H2C	7	43	28	47	50	44	50	48
4	25	90	95	100	90	93	100	90
D1A	17	68	66	67	75	70	45	51
S36	4	60	57	60	0	0	60	6
H17C	12	110	110	110	110	120	120	120
F1A	17	50	55	60	62	62	49	69
H7C	5	58	32	52	73	62	35	44
S9	22	84	86	86	86	86	94	85
H11B	10	45	32	52	47	54	54	62
H10A	29	27	62	48	71	63	27	—
H10B	37	44	75	75	80	72	100	57

TABLE 4

Effect of Homobiotin upon the Growth of *Staphylococcus* sp. H17C.

Homobiotin (μg./10 ml.)	Optical density x 100 (41 hrs.)	
	With biotin (2 mμg./10 ml.)	Without biotin
0	110	15
0.2	110	5
2.0	110	32
20	110	95
200	110	130
2000	110	130

TABLE 5  
Comparative Activity of Homobiotin for  
Different *Staphylococcus* Strains.

Culture	Homobiotin necessary for 1/2 maximum growth ( $\mu\text{g.}/10\text{ ml.}$ )
H9A	1
H13A	0.3-1
D2A	0.3
H17C	1-3
S19	0.3-1
H3A	0.4-0.8
H2C	0.4
H10A*	10-30

\* Culture H10A also requires unusually large amounts of biotin for 1/2 maximum growth.

TABLE 6  
Comparative Activities of Biotin and Biocytin.

Culture	Quantities required for 1/2 max. growth ( $\text{m}\mu\text{g.}/10\text{ ml.}$ )		Biocytin Biotin
	Biotin	Biocytin	
H3A	.03 — .05	.05 — 0.1	1.9
D2A	.05	0.1	2.0
H10A	5 — 10	10 — 30	1.4
	10 — 30		

## REFERENCES

- BELCHER, M. RUTH, and LICHTSTEIN, HERMAN C. — 1949 — Growth promotion and antibiotic effect of homobiotin and norbiotin. *J. Bact.*, **58**, 579-583.
- EVANS, J. B. — 1948 — Studies of staphylococci with special reference to the coagulase-positive types. *J. Bact.*, **55**, 793-800.
- EVANS, JAMES B., and NIVEN, C. F. JR. — 1950 — A comparative study of known food-poisoning staphylococci and related varieties. *J. Bact.*, **59**, 545-550.
- EVANS, JAMES B., BUETTNER, L. G., and NIVEN, C. F., JR. — 1950 — Evaluation of coagulase test in the study of staphylococci associated with food poisoning. *J. Bact.*, **60**, 481-484.
- GOLDBERG, M. W., STERNBACH, L. H., KAISER, S., HEINEMAN, S. D., SCHEINER, J., and RUBIN, S. H. — 1947 — Antibiotin effect of some biotin homologs. *Arch. Biochemistry*, **14**, 480-482.
- GRETTLER, ALBERT C. — 1950 — A study of the vitamin nutrition of staphylococci with special reference to biotin requirements. Thesis, Department of Bacteriology and Parasitology, University of Chicago. December, 1950.
- GRETTLER, ALBERT C., and EVANS, JAMES B. — 1951 — Vitamin requirements of staphylococci. *Bact. Proc., Soc. Am. Bact.*, 1951.
- HUTNER, S. H., and BJERKNES, C. A. — 1948 — Volatile preservatives for culture media. *Proc. Soc. Exptl. Biol. Med.*, **67**, 393-397.
- RUBIN, S. H., DREKTER, L., and MAYER, E. H. — 1945 — Biological activity of synthetic d,l-desfthiobiotin. *Proc. Soc. Exptl. Biol. Med.*, **58**, 352-356.
- RUBIN, SAUL H., and SCHEINER, JACOB — 1949 — Antibiotic effect of homologs of biotin and biotin sulfone. *Arch. Biochemistry*, **23**, 400-410.
- STOKES, JACOB L., and GUNNESS, MARION — 1945 — Microbiological activity of synthetic biotin, its optical isomers, and related compounds. *J. Biol. Chem.*, **157**, 121-126.
- WRIGHT, L. D., CRESSON, E. L., SKEGGS, H. R., WOOD, T. R., PECK, R. L., WOLF, D. E., and FOLKERS, K. — 1950a — Biocytin, a naturally-occurring complex of biotin. *J. Am. Chem. Soc.*, **72**, 1048.
- WRIGHT, LEMUEL D., CRESSON, EMELON L., VALENTICK, KATHERINE and SKEGGS, HELEN R. — 1950b — Microbiological characterization of biocytin. *Fed. Proc.*, **9**, 374.