

Quantum Stochastic Processes, Quantum Iterated Function Systems and Entropy

A. Baraviera; C. F. Lardizabal

A. O. Lopes and M. Terra Cunha

Abstract. We describe some basic results for Quantum Stochastic Processes and present some new results about a certain class of processes which are associated to Quantum Iterated Function Systems (QIFS). We discuss questions related to the Markov property and we present a definition of entropy which is induced by a QIFS. This definition is a natural generalization of the Shannon-Kolmogorov entropy from Ergodic Theory.

1. Introduction

We review and discuss some main properties of Quantum Stochastic Processes (see [6] [18] [20]) and present some new results about a certain class of processes which are associated to a Quantum Iterated Function System (QIFS). The concept of QIFS was introduced in the work [14], and it is a natural object in Quantum Information Theory.

We also present a definition of entropy which is suitable for the QIFS. This definition is a natural generalization of the Shannon-Kolmogorov entropy of Ergodic Theory. We describe a parallel between the classical Kolmogorov entropy and the one we present here, which is different from the one seen in [1].

The present definition of entropy is obtained by adapting the reasoning described in [5], [12] and [13] to the setting we present in this work. The main idea is to define this concept via the Ruelle operator and to avoid the use of partitions. Using this definition one can consider maximal pressure

Supported in part by CAPES and CNPq.

density states. This formulation can be seen as a mini-max problem (see [5] [12] [13]). In [1] it is described some applications of the pressure problem.

Section 2 introduces basic notations and section 3 describes QSPs following [18]; section 4 and 5 describes Quantum Iterated Function Systems, following [14]. Section 6 is a brief digression on the Chapman-Kolmogorov equation and probability amplitudes. Section 7 defines probabilities measures and quantum stochastic processes induced by QIFS. Section 8 gives a definition of entropy induced by a QIFS and we make a few remarks on the variational problem of pressure.

Our work is inspired by results presented in [14] and [19]. We would like to thank these authors for supplying us with the corresponding references. Some other references related to the topic described here are [1] [2] [3] [4] [19].

This work is part of the thesis dissertation of C. F. Lardizabal in Prog. Pos-Grad. Mat. UFRGS (Brazil) [10].

2. Notations

We recall some basic notation which is used in Quantum Computing. For a comprehensive introduction to the subject, see [16]. Let \mathcal{H}_N be a Hilbert space of finite dimension N . A **state** in a quantum system is described by $|\psi\rangle \in \mathcal{H}_N$. Such states are normalized, so we have $\langle\psi|\psi\rangle = 1$. For any phase α , we identify the elements $|\psi'\rangle = e^{i\alpha}|\psi\rangle$ and $|\psi\rangle$, so we get the space of pure states, denoted by \mathcal{P}_N . Topologically, it is the complex projective space \mathbb{CP}^{N-1} with the Fubini-Study metric, given by $D_{FS}(|\phi\rangle, |\psi\rangle) := \arccos|\langle\phi|\psi\rangle|$.

A **qubit** is a unit vector in a complex vector space of dimension 2

$$|\psi\rangle = \alpha|0\rangle + \beta|1\rangle,$$

where $|\alpha|^2 + |\beta|^2 = 1$. We can rewrite such equation as

$$|\psi\rangle = e^{i\gamma}(\cos\frac{\theta}{2}|0\rangle + e^{i\phi}\sin\frac{\theta}{2}|1\rangle),$$

where θ, ϕ, γ are real numbers. As we are in projective space, the factor $e^{i\gamma}$ can be ignored, so we can write

$$|\psi\rangle = \cos\frac{\theta}{2}|0\rangle + e^{i\phi}\sin\frac{\theta}{2}|1\rangle$$

The numbers θ and ϕ define a point on the unit sphere, the Bloch sphere, which gives us an easy way to visualize the state of a qubit.

Denote by ρ^* the adjoint of $\rho : \mathcal{H}_N \rightarrow \mathcal{H}_N$. We say that $\rho : \mathcal{H}_N \rightarrow \mathcal{H}_N$ is hermitian if $\rho = \rho^*$. We say that a hermitian operator $P : \mathcal{H}_N \rightarrow \mathcal{H}_N$ is positive, denoting such fact by $P \geq 0$, if $\langle Pv, v \rangle \geq 0$, $\forall v \in \mathcal{H}_N$.

Definition 1. A **density operator** (or *density matrix*) is an operator ρ acting on \mathcal{H}_N , with $\rho = \rho^*$, $\rho \geq 0$ and $\text{tr} \rho = 1$. Denote by \mathcal{M}_N the space of density operators.

If $|\psi\rangle \in \mathcal{H}_N$ is a state, denote its **associated projection** by $|\psi\rangle\langle\psi|$ (which defines an associated density matrix). We denote by $\{|0\rangle, \dots, |N-1\rangle\}$ the canonical orthonormal basis for \mathcal{H}_N . A density operator ρ can always be written as

$$\rho = \sum_{i=1}^k p_i |\psi_i\rangle\langle\psi_i| \quad (1)$$

where the p_i are positive numbers with $\sum_i p_i = 1$, and $|\psi_i\rangle$, $i = 1, 2, \dots, N-1$, is an orthonormal basis. So the $|\psi_i\rangle$ are eigenvectors of ρ with the p_i as their respective eigenvalues.

If a density operator satisfies $\text{tr}(\rho^2) = 1$ then such operator is a projection and we say that ρ represents a **pure state**. If ρ is a nontrivial convex combination of projections then we say ρ represents a **mixed state**; in this case we have $\text{tr}(\rho^2) < 1$. Also an operator is a density operator if and only if its trace equals 1 and if it is positive.

3. A description of quantum stochastic process

In this section the definitions and examples were taken from [18], where it is presented a definition of quantum stochastic process. We briefly describe some of the results obtained in that work.

Definition 2. A *state space* is a pair (V, K) , where

- (1) V is a real Banach space with norm $\|\cdot\|$.
- (2) K is a closed cone in V .
- (3) If $u, v \in K$ then $\|u\| + \|v\| = \|u + v\|$
- (4) If $u \in V$ and $\epsilon > 0$ then there are $u_1, u_2 \in K$ such that $u = u_1 - u_2$ and $\|u_1\| + \|u_2\| < \|u\| + \epsilon$.

Definition 3. If (V, K) is a state space then there is a unique positive linear functional $\tau : V \rightarrow \mathbb{R}$ such that $\tau(u) = \|u\|$ if $u \in K$, and $\tau(u) \leq \|u\|$ if $u \in V$. We say that $u \in K$ is a **state** if $\tau(u) = 1$.

Example 1. Let \mathcal{H} be a finite dimensional Hilbert space and let V be the space of hermitian operators in \mathcal{H} . Let K be the set of positive operators in V . In this case we have $\tau(B) = \text{tr}(B)$ for all B operator in V .

◇

Definition 4. A **phase space** is a measurable space (Ω, Σ) where Ω represents the set of all possible results for a measurement and Σ is a σ -algebra of subsets of Ω .

Let V^* be the dual space of V . We introduce a partial order on V^* by defining $\phi \geq \psi$ if $\phi(u) \geq \psi(u)$, for all $u \in K$.

Definition 5. An **effect** is a mapping $\phi \in V^*$ such that $0 \leq \phi \leq \tau$. We denote the space of effects by $\mathcal{E} \subset V^*$.

Definition 6. We say that $x : \Sigma \rightarrow \mathcal{E}$ is an **observable** if x is a measure taking values on the space of effects, such that $x(\Omega) = \tau$.

If $E \in \Sigma$, $u \in K$ and $\tau(u) = 1$ then $x(E)u$ can be interpreted as the probability that the result of the measurement of the physical quantity represented by x , prepared in the state u , belongs to the set E . In the case of quantum mechanics in Hilbert space, effects can be identified with bounded operators A such that $0 \leq A \leq 1$ by the formula $\phi_A(W) = \text{tr}(AW)$.

Definition 7. An **operation** is a positive linear operator $T : V \rightarrow V$ satisfying $0 \leq \tau(Tu) \leq \tau(u)$ for all $u \in K$. The space of operations will be denoted by \mathcal{O} .

Definition 8. An **operator valued measure**, or an **OVM** on a phase space is a map $\mathcal{I} : \Sigma \rightarrow \mathcal{O}$ such that if $\{E_n\}$ is a sequence of disjoint sets in Σ , then $\mathcal{I}(\cup E_n) = \sum \mathcal{I}(E_n)$.

Definition 9. Let $\mathcal{I} : \Sigma \rightarrow \mathcal{O}$ be an OVM, then we say that \mathcal{I} is an **instrument** if

$$\tau(\mathcal{I}(\Omega)u) = \tau(u), \forall u \in V. \quad (2)$$

We interpret such notion in the following way. Let \mathcal{I} be an instrument, $E \in \Sigma$, $u \in K$. If u is the state of the system before the measurement and if \mathcal{I} determines a value in E then the resulting state is given by

$$\frac{\mathcal{I}(E)u}{\tau(\mathcal{I}(E)u)} \quad (3)$$

Note that for each instrument \mathcal{I} , there is a unique observable $x_{\mathcal{I}} : \Sigma \rightarrow \mathcal{E}$ such that $\tau(\mathcal{I}(E)u) = x_{\mathcal{I}}(E)u$, $E \in \Sigma$, $u \in K$. Also, it is possible that two instruments correspond to the same observable [18].

The following are examples of instruments:

Example 2. Let \mathcal{H} be a Hilbert space, and let $\mathcal{F}(\mathcal{H})$ be the space of hermitian operators A in \mathcal{H} such that

$$\sum_{k \in \mathbb{N}} \langle e_k, A e_k \rangle < \infty$$

and have the same value in any orthonormal base $\{e_k\}_{k \in \mathbb{N}}$ for \mathcal{H} . Let $\Omega = \{1, \dots, N\}$, or $\Omega = \mathbb{N}$, let $\{P_i\}_{i \in \Omega}$ be a family of orthogonal projections such that $\sum_i P_i = I$. Define

$$\mathcal{I} : \Sigma \rightarrow \mathcal{O}$$

$$x_{\mathcal{I}} : \Sigma \rightarrow \mathcal{E}$$

as

$$\mathcal{I}(E)\rho := \sum_{i \in E} P_i \rho P_i, \quad (4)$$

$$x_{\mathcal{I}}(E)\rho := \sum_{i \in E} \tau(P_i \rho), \quad (5)$$

for all $E \subset \Omega$ and $\rho \in \mathcal{F}(\mathcal{H})$.

◇

Example 3. Let \mathcal{H} be a Hilbert space, Ω a topological space, Σ a σ -algebra for Ω and m a measure on (Ω, Σ) . Let $\{P_a\}_{a \in \Omega}$ be a family of projections on \mathcal{H} , such that the mapping $a \rightarrow P_a$ is strongly continuous and $\int_{\Omega} P_a dm(a) = I$. Then define

$$\mathcal{I} : \Sigma \rightarrow \mathcal{O}$$

$$x_{\mathcal{I}} : \Sigma \rightarrow \mathcal{E}$$

as

$$\mathcal{I}(E)\rho := \int_E P_a \rho P_a dm(a) \quad (6)$$

$$x_{\mathcal{I}}(E)\rho := \int_E \tau(P_a \rho) dm(a), \quad (7)$$

for all $E \subset \Omega$ e $\rho \in \mathcal{F}(\mathcal{H})$.

◇

Example 4. Let X be a locally compact Hausdorff space, V the space of the countably additive functions on the Borel σ -algebra $\mathcal{B}(X)$ for X endowed with the norm of total variation. Let K be the set of nonnegative measures on V . Let $(\Omega, \Sigma) = (X, \mathcal{B}(X))$. Then

$$\mathcal{I}(E)\mu(A) = \mu(A \cap E), \quad (8)$$

for $\mu \in V$, $A, E \in \Sigma$ is an instrument, called the **sharp classical measurement** and the corresponding observable is

$$x_{\mathcal{I}}(E)\mu = \mu(E) \quad (9)$$

◇

Definition 10. *Following [18], a **Quantum Stochastic Process, QSP**, is an arbitrary family of instruments $\{\mathcal{I}_t\}_{t \in \mathcal{J}}$. Let $\mathcal{J} = \mathbb{Z}$ or $\mathcal{J} = \mathbb{R}$ for discrete or continuous time, respectively.*

The **finite dimensional distributions** of the process are measures $\mu_{t_0, \dots, t_{n-1}}^u$ defined in $(\Omega^n, B(\Omega^n))$ as being the natural extensions of the functions given by

$$\mu_{t_0, \dots, t_{n-1}}^u(E_0 \times \dots \times E_{n-1}) = \tau((\mathcal{I}_{t_{n-1}}(E_{n-1}) \circ \mathcal{I}_{t_{n-2}}(E_{n-2}) \circ \dots \circ \mathcal{I}_{t_0}(E_0))u) \quad (10)$$

where $n \in \mathbb{N}$, $t_0 \leq \dots \leq t_{n-1}$, $t_i \in \mathcal{J}$, $u \in V$ and $E_0, \dots, E_{n-1} \in \Sigma$. The meaning of such expression is the following: $\mu_{t_0, \dots, t_{n-1}}^u(E_0 \times \dots \times E_{n-1})$ is the joint probability that successive measurements of the system by the instruments $\mathcal{I}_0, \dots, \mathcal{I}_{n-1}$ in the moments t_0, \dots, t_{n-1} produce values in E_0, \dots, E_{n-1} , when the pre-measurement state is u .

A **probability transition** is a function $P : \Omega \times \Sigma \rightarrow \mathbb{R}$ such that $P(\cdot, E)$ is measurable for all $E \in \Sigma$ and $P(x, \cdot)$ is a probability measure for all $x \in \Omega$.

Definition 11. *We say that a QSP is **Markov** if there exists a family of probability transitions $\{P_{s,t}\}_{s < t}$ such that*

$$\begin{aligned} & \mu_{t_0, \dots, t_{n-1}}^u(E_0 \times \dots \times E_{n-1}) \\ &= \int_{E_0} \int_{E_1} \dots \int_{E_n} P_{t_{n-1}, t_n}(y_{n-1}, dy_n) \dots P_{t_0, t_1}(y_0, dy_1) \mu_{t_0}^u(dy_0) \end{aligned} \quad (11)$$

for all $t_0 < \dots < t_n$, $t_i \in \mathcal{J}$, $u \in V$, $E_0, \dots, E_n \in \Sigma$. A Markov QSP is **homogeneous** if the probability transitions $P_{s,t}$ depend only on the difference $t - s$.

Remark In contrast with the classic theory of stochastic processes, the probability transitions of a Markov QSP do not satisfy in general the Chapman-Kolmogorov equation.

◇

Definition 12. Let \mathcal{I} be an instrument. Assume that between the measurements the system evolves and its evolution is described by a group $\{T_t\}_{t \in \mathcal{J}}$ of isometric automorphisms of V . Then define the QSP $\{\mathcal{I}_t\}_{t \in \mathcal{J}}$, where

$$\mathcal{I}_t(E) = T_t^{-1} \circ \mathcal{I}(E) \circ T_t \quad (12)$$

is called a **transformed instrument**. For simplicity, we can choose $\mathcal{J} = \mathbb{Z}$ so $T_n = T^n$ and we denote such process by $\mathcal{C}(T, \mathcal{I})$.

Now we show an example of a Markov QSP.

Example 5. Let \mathcal{I} be the instrument given in example 4 and let $\Theta : X \rightarrow X$ be a measurable map. Then Θ generates an automorphism $T_\Theta : V \rightarrow V$ by

$$T_\Theta(\mu)(A) = \mu(\Theta^{-1}(A)), \quad \mu \in V, A \in B(X) \quad (13)$$

Then we can show that $\mathcal{C}(T_\Theta, \mathcal{I})$ is a homogeneous Markov QSP and its transition probability is given by

$$P(x, E) = \mathcal{X}_E(\Theta x), \quad x \in X, E \in B(X) \quad (14)$$

◇

4. Quantum IFS

This section follows [14]. We begin with a few definitions.

Definition 13. Let $G_i : \mathcal{M}_N \rightarrow \mathcal{M}_N$, $p_i : \mathcal{M}_N \rightarrow [0, 1]$, $i = 1, \dots, k$ and such that $\sum_i p_i(\rho) = 1$. We call

$$\mathcal{F}_N = \{\mathcal{M}_N, G_i, p_i : i = 1, \dots, k\} \quad (15)$$

a **Quantum Iterated Function System (QIFS)**.

Definition 14. A QIFS is **homogeneous** if p_i and $G_i p_i$ are affine mappings, $i = 1, \dots, k$.

Suppose that the QIFS considered is such that there are V_i and W_i linear maps, $i = 1, \dots, k$, with $\sum_{i=1}^k W_i^* W_i = I$ such that

$$G_i(\rho) = \frac{V_i \rho V_i^*}{\text{tr}(V_i \rho V_i^*)} \quad (16)$$

and

$$p_i(\rho) = \text{tr}(W_i \rho W_i^*) \quad (17)$$

Then we have that a QIFS is homogeneous if $V_i = W_i$, $i = 1, \dots, k$. Now we can define a Markov operator $P : \mathcal{M}(\mathcal{M}_N) \rightarrow \mathcal{M}(\mathcal{M}_N)$,

$$(P\mu)(B) = \sum_{i=1}^k \int_{G_i^{-1}(B)} p_i(\rho) d\mu(\rho),$$

where $\mathcal{M}(\mathcal{M}_N)$ denotes the space of probability measure over \mathcal{M}_N . We also define $\Lambda : \mathcal{M}_N \rightarrow \mathcal{M}_N$,

$$\Lambda(\rho) := \sum_{i=1}^k p_i(\rho) G_i(\rho)$$

If the QIFS considered is homogeneous then

$$\Lambda(\rho) = \sum_i V_i \rho V_i^* \quad (18)$$

We say that $\rho \in \mathcal{M}_N$ is the **integral** of a mapping $f : \mathcal{M}_N \rightarrow \mathcal{M}_N$, denoted by

$$\rho := \int_{\mathcal{M}_N} f d\mu$$

if

$$l(\rho) = \int_{\mathcal{M}_N} l \circ f d\mu,$$

for all $l \in \mathcal{M}_N^*$.

Theorem 1. *A mixed state ρ_0 is Λ -invariant, if and only if,*

$$\rho_0 = \int_{\mathcal{M}_N} \rho d\mu(\rho), \quad (19)$$

for some P -invariant measure μ .

For the proof, see [14], [19].

In order to define hyperbolic QIFS, we have to specify a distance on the space of mixed states. The following are a few possibilities:

$$\begin{aligned} D_1(\rho_1, \rho_2) &= \sqrt{\text{tr}[(\rho_1 - \rho_2)^2]} \\ D_2(\rho_1, \rho_2) &= \text{tr} \sqrt{(\rho_1 - \rho_2)^2} \\ D_3(\rho_1, \rho_2) &= \sqrt{2\{1 - \text{tr}[(\rho_1^{1/2} \rho_2 \rho_1^{1/2})^{1/2}]\}} \end{aligned}$$

Such metrics generate the same topology on \mathcal{M}_N . Considering the space of mixed states with one of those metrics we can make the following definition.

Definition 15. *We say a QIFS is **hyperbolic** if the quantum maps G_i are contractions with respect to one of the distances on \mathcal{M}_N and if the maps p_i are Hölder-continuous and positive.*

Proposition 1. [14] [19] *If a QIFS (15) is homogeneous and hyperbolic then the associated Markov operator admits a unique invariant measure μ . Such invariant measure determines a unique Λ -invariant state $\rho \in \mathcal{M}_N$, given by (19).*

5. Examples of QIFS

Example 6. $\Omega = \mathcal{M}_N$, $k = 2$, $p_1 = p_2 = 1/2$, $G_1(\rho) = U_1 \rho U_1^*$, $G_2(\rho) = U_2 \rho U_2^*$. The normalized identity matrix $\rho_* = I/N$ is Λ -invariant, for any choice of unitary U_1 and U_2 . Note that we can write

$$\rho_* = \int_{\mathcal{M}_N} \rho d\mu(\rho)$$

where the measure μ , uniformly distributed over \mathcal{P}_N , is P -invariant. \diamond

Example 7. Let $\Omega = \mathcal{M}_N$, $k = 2$, $p_1 = p_2 = 1/2$, $G_1(\rho) = (\rho + 2\rho_1)/3$, $G_2(\rho) = (\rho + 2\rho_2)/3$, where we choose the projectors $\rho_1 = |1\rangle\langle 1|$ and $\rho_2 = |2\rangle\langle 2|$ so that they are orthogonal. Since G_1 and G_2 are contractions with Lipschitz constant equal to $1/3$, this QIFS is hyperbolic and so there is a unique invariant measure. \diamond

Recall that a mapping Λ is **completely positive** (CP), if $\Lambda \otimes I$ is positive for any extension of the original Hilbert space $\mathcal{H}_N \rightarrow \mathcal{H}_N \otimes \mathcal{H}_E$. We know that every trace preserving CP map can be represented (in a nonunique way) in the Stinespring-Kraus form

$$\Lambda(\rho) = \sum_{j=1}^k V_j \rho V_j^*, \quad \sum_{j=1}^k V_j^* V_j = 1,$$

where the V_j are linear operators. Besides, if $\sum_{j=1}^k V_j V_j^* = I$ then $\Lambda(I/N) = I/N$ and Λ will be called **unital**. This is the case if each of the V_j is normal, that is, if $V_j V_j^* = V_j^* V_j$. Note that by writing $G_i(\rho) = U_i \rho U_i^*$, we have that example 6 is contained in this class of QIFS. We call such QIFS **unitary**. For a unitary QIFS we have that ρ_* is an invariant state for Λ_U and also that δ_{ρ_*} is invariant for the Markov operator P_U induced by this QIFS.

Definition 16. We say that unitary matrices of same dimension are **common block diagonal** if they are block diagonal in the same base and with the same blocks.

The proof of the following lemma is presented in [14].

Proposition 2. Assume that p_i , $i = 1, \dots, k$ are strictly positive. The the maximally mixed state ρ_* is the unique invariant state for the operator Λ_U if and only if the unitary operators U_i , $i = 1, \dots, k$ are not common block diagonal.

Example 8. Let $\Omega = \mathcal{P}_2$, $U_1 = I$, $U_2 = \sigma_1$, $U_3 = \sigma_2$, $U_4 = \sigma_3$, $p_1 = 1 - p$, $p_2 = p_3 = p_4 = p/3 > 0$, where $\sigma_1, \sigma_2, \sigma_3$ are the Pauli matrices. Since such matrices are not common block diagonal the maximally mixed state ρ_* is the unique invariant state for the mapping below, called a quantum depolarizing channel [14]:

$$\Lambda_U(\rho) = \sum p_i U_i \rho U_i^* = (1 - p)\rho + \frac{p}{3}(\sigma_1 \rho \sigma_1 + \sigma_2 \rho \sigma_2 + \sigma_3 \rho \sigma_3).$$

◇

Example 9. Let $\Omega = \mathcal{P}_2$, $p_1 = 1 - p$, $p_2 = p$,

$$U_1 = \exp(-iH_0T/\hbar),$$

$$U_2 = \exp(-\frac{i}{\hbar}(H_0T + \int_0^T V(t)dt))$$

where $V(t) = V(t + T)$. The maximally mixed state $\rho_* = I/2$ is an invariant state for the operator Λ_U corresponding to this QIFS. For a generic perturbation V , matrices U_1 and U_2 are not common block diagonal so ρ_* is the unique invariant state for Λ_U .

◇

6. On certain probability and amplitude calculations

We begin with a brief digression on the Chapman-Kolmogorov equation. Let $X = \{X_n\}$ be a sequence of measurable functions. Suppose that

$$P(X_{n+1} = j | X_n = i) = P(X_1 = j | X_0 = i)$$

for all n, i, j . Suppose that X takes values on a finite set S . Define the matrix $P = (p_{ij})$ of order $|S|$, with entries

$$p_{ij} = P(X_{n+1} = j | X_n = i)$$

Define the matrix of n transitions $P_n = (p_{ij}(n))$, where

$$p_{ij}(n) = P(X_{m+n} = j | X_m = i)$$

Also suppose that it is a Markov chain, that is

$$P(X_n = x_n | X_0 = x_0, X_1 = x_1, \dots, X_{n-1} = x_{n-1}) = \quad (20)$$

$$P(X_n = x_n | X_{n-1} = x_{n-1})$$

for all $n \geq 1$, and $x_0, \dots, x_n \in S$.

By using the fact that for any events A_1, A_2, A_3 , we have

$$P(A_1 \cap A_2 | A_3) = P(A_1 | A_2 \cap A_3)P(A_2 | A_3) \quad (21)$$

we can write

$$\begin{aligned} p_{ij}(m+n) &= P(X_{m+n} = j | X_0 = i) = \sum_k P(X_{m+n} = j, X_m = k | X_0 = i) \\ &= \sum_k P(X_{m+n} = j | X_m = k) P(X_m = k | X_0 = i) \end{aligned} \quad (22)$$

So

$$p_{ij}(m+n) = \sum_k p_{ik}(m) p_{kj}(n) \quad (23)$$

which is the Chapman-Kolmogorov equation. We are interested in studying quantum stochastic processes and in obtaining an adequate definition to what we will call a Markov quantum stochastic process. First we recall that in the previous section we have presented a description [18] of Markov QSP in which the Chapman-Kolmogorov do not hold in general. This fact can be seen as the general rule for quantum processes (but see [8] for different settings).

In algebraic terms, we can argue that the deduction of (23) above is not valid for quantum processes because of equation (21). Since we have to take in consideration the interference between measurements, the problem of understanding how probability measures work in a quantum setting is a basic question. In quantum mechanics we could in principle consider a probability space (Ω, Λ, μ) such as in classic measure theory. However, we have that Λ is a σ -algebra and μ is a measure on Λ only when we are restricted to a single measurement. When we perform several measurements interference effects occur and so we are no longer considering a problem on classic probability [7]. Results of more general nature are presented in [9].

We can think that interference occurs because, in contrast to classic probability measures, which can be quite arbitrary, quantum probability measures are obtained in a very specific way. In quantum mechanics we have an amplitude function $a : \Omega \rightarrow \mathbb{C}$, and if $B \in \Lambda$, we define the **amplitude** of B as

$$A(B) = \sum_{\omega \in B} a(\omega) \quad (24)$$

and we define the probability that B occurs as

$$\mu(B) = |A(B)|^2 \quad (25)$$

Let us describe a few more details on this point. For more on the subject, see for instance [7]. Let Ω be a nonempty set and let $a : \Omega \rightarrow \mathbb{C}$. We say that $\omega \in \Omega$ is a **sample point** and the map a is a **probability amplitude**, and (Ω, f) is called a **quantum probability space**. A set $A \subset \Omega$ is **summable**

if $\sum_{\omega \in \Omega} |a(\omega)|^2 < \infty$ and we denote the collection of summable sets by Σ_0 . Now define $A : \Sigma_0 \rightarrow \mathbb{C}$ as $A(\emptyset) = 0$ and

$$A(B) := \sum_{\omega \in \Omega} a(\omega) \quad (26)$$

We say that $A(B)$ is the **amplitude** of B . Now define

$$A(B_1|B_2) := \frac{A(B_1 \cap B_2)}{A(B_2)} \quad (27)$$

if $A(B_2) \neq 0$ and equal to zero, otherwise. In the case that $A(B_2) \neq 0$, we have that $A(\cdot|B_2)$ is a complex measure on $P(\Omega)$, with $A(\Omega|B_2) = 1$. We say that $A(B_1|B_2)$ is the **conditional amplitude** of B_1 , given B_2 . Note that $A(B) = 0$ does not imply $A(B \cap C) = 0$ [7]. Because of that, formulas of the kind $A(B \cap C) = A(B)A(C|B)$ might not be true when $A(B) = 0$. However, when the conditioning sets have a nonzero amplitude, we have the formula

$$A(B_1 \cap \dots \cap B_n) = A(B_1)A(B_2|B_1)A(B_3|B_1 \cap B_2) \dots A(B_n|B_1 \cap \dots \cap B_{n-1}) \quad (28)$$

which is the amplitude counterpart for equation (21). Define the matrix $A = (a_{ij})$, where $a_{ij} = A(X_{n+1} = j|X_n = i)$. Now suppose that the chain $\{X_n\}_{n \in \mathbb{N}}$ is **quantum Markov**, that is,

$$A(X_n = x_n|X_0 = x_0, X_1 = x_1, \dots, X_{n-1} = x_{n-1}) = A(X_n = x_n|X_{n-1} = x_{n-1}) \quad (29)$$

for all $n \geq 1$, $x_0, \dots, x_n \in S$. So in a way which is similar to what we did for probabilities, define the matrix of n transitions $A_n = (a_{ij}(n))$, where $a_{ij}(n) = A(X_{m+n} = j|X_m = i)$ and we get

$$a_{ij}(m+n) = \sum_k a_{ik}(m)a_{kj}(n) \quad (30)$$

so we have that $A_{m+n} = A_m A_n$ and $A_n = A^n$.

◇

7. Probability measures induced by QIFS

In this section we present some new results. Consider a Hilbert space \mathcal{H} of dimension $N = 2$. Let $q_1, q_2 \in \mathbb{R}$ and also

$$V_1 = \begin{pmatrix} \sqrt{p_{11}} & \sqrt{p_{12}} \\ 0 & 0 \end{pmatrix}, \quad V_2 = \begin{pmatrix} 0 & 0 \\ \sqrt{p_{21}} & \sqrt{p_{22}} \end{pmatrix}, \quad \rho = \begin{pmatrix} \rho_1 & \rho_2 \\ \rho_3 & \rho_4 \end{pmatrix} \quad (31)$$

We would like to obtain the fixed points for

$$\mathcal{L}(\rho) = q_1 V_1 \rho V_1^* + q_2 V_2 \rho V_2^*$$

Then

$$q_1 V_1 \rho V_1^* + q_2 V_2 \rho V_2^* = \rho \quad (32)$$

implies

$$q_1 \left[(\sqrt{p_{11}}\rho_1 + \sqrt{p_{12}}\rho_3)\sqrt{p_{11}} + (\sqrt{p_{11}}\rho_2 + \sqrt{p_{12}}\rho_4)\sqrt{p_{12}} \right] = \rho_1$$

$$q_2 \left[(\sqrt{p_{21}}\rho_1 + \sqrt{p_{22}}\rho_3)\sqrt{p_{21}} + (\sqrt{p_{21}}\rho_2 + \sqrt{p_{22}}\rho_4)\sqrt{p_{22}} \right] = \rho_4$$

And (32) also implies that $\rho_2 = \rho_3 = 0$, so we rewrite the system as

$$q_1 \left[\sqrt{p_{11}}\rho_1\sqrt{p_{11}} + \sqrt{p_{12}}\rho_4\sqrt{p_{12}} \right] = \rho_1$$

$$q_2 \left[\sqrt{p_{21}}\rho_1\sqrt{p_{21}} + \sqrt{p_{22}}\rho_4\sqrt{p_{22}} \right] = \rho_4$$

or

$$a\rho_1 + f\rho_4 = \rho_1 \quad (33)$$

$$g\rho_1 + h\rho_4 = \rho_4 \quad (34)$$

where

$$a = q_1 p_{11}, \quad f = q_1 p_{12}, \quad g = q_2 p_{21}, \quad h = q_2 p_{22}$$

We also get that

$$\rho_1 = \frac{f}{1-a}\rho_4$$

$$\rho_1 = \frac{1-h}{g}\rho_4$$

which is a restriction on the q_i , namely

$$\frac{f}{1-a} = \frac{1-h}{g}$$

Therefore the solution of (33) and (34) is

$$\rho = \rho_4 \begin{pmatrix} \frac{f}{1-a} & 0 \\ 0 & 1 \end{pmatrix} = \rho_4 \begin{pmatrix} \frac{1-h}{g} & 0 \\ 0 & 1 \end{pmatrix}$$

But $\rho_1 + \rho_4 = 1$ implies

$$\rho = \begin{pmatrix} \frac{q_1 p_{12}}{q_1 p_{12} - q_1 p_{11} + 1} & 0 \\ 0 & \frac{1 - q_1 p_{11}}{q_1 p_{12} - q_1 p_{11} + 1} \end{pmatrix} = \begin{pmatrix} \frac{1 - q_2 p_{22}}{1 - q_2 p_{22} + q_2 p_{21}} & 0 \\ 0 & \frac{q_2 p_{21}}{1 - q_2 p_{22} + q_2 p_{21}} \end{pmatrix} \quad (35)$$

Now assume that

$$P = \begin{pmatrix} p_{11} & p_{12} \\ p_{21} & p_{22} \end{pmatrix}$$

is column stochastic, that is, we have $p_{ij} \geq 0$ and $\sum_i p_{ij} = 1$, $i, j = 1, 2$. Let π be such that $P\pi = \pi$. Such π is given by

$$\pi = \left(\frac{p_{12}}{p_{12} - p_{11} + 1}, \frac{1 - p_{11}}{p_{12} - p_{11} + 1} \right) \quad (36)$$

Compare (36) with (35). Then fix $q_1 = q_2 = 1$, so we get that the nonzero entries of ρ are equal to the entries of π . Such a choice for the q_i is unique. In fact, comparing the (i, i) -th entry of ρ with the i -th coordinate of π , we see that if there exists q'_i which make ρ and π equal (i.e., the diagonal entries of ρ correspond to the entries of π), then

$$\frac{q_1 p_{12}}{q_1 p_{12} - q_1 p_{11} + 1} = \frac{q'_1 p_{12}}{q'_1 p_{12} - q'_1 p_{11} + 1},$$

which implies

$$\begin{aligned} q_1(q'_1 p_{12} - q'_1 p_{11} + 1) &= q'_1(q_1 p_{12} - q_1 p_{11} + 1) \\ \Rightarrow q_1 q'_1 p_{12} - q_1 q'_1 p_{11} + q_1 &= q_1 q'_1 p_{12} - q_1 q'_1 p_{11} + q'_1 \end{aligned}$$

and when we cancel terms we get $q_1 = q'_1$. In a similar way

$$\frac{1 - q_2 p_{22}}{1 - q_2 p_{22} + q_2 p_{21}} = \frac{1 - q'_2 p_{22}}{1 - q'_2 p_{22} + q'_2 p_{21}}$$

implies

$$\begin{aligned} (1 - q_2 p_{22})(1 - q'_2 p_{22} + q'_2 p_{21}) &= (1 - q'_2 p_{22})(1 - q_2 p_{22} + q_2 p_{21}) \\ \Rightarrow 1 - q'_2 p_{22} + q'_2 p_{21} - q_2 p_{22} + q_2 q'_2 p_{22}^2 - q_2 q'_2 p_{22} p_{21} \\ &= 1 - q_2 p_{22} + q_2 p_{21} - q'_2 p_{22} + q_2 q'_2 p_{22}^2 - q_2 q'_2 p_{22} p_{21} \end{aligned}$$

Then we get

$$q'_2 p_{21} = q_2 p_{21} \Rightarrow q'_2 = q_2$$

and therefore the choice for q_1 and q_2 is unique.

◇

Consider a homogeneous QIFS $\mathcal{F} = \{\mathcal{M}_N, F_i, p_i\}_{i=1, \dots, k}$, where

$$F_i(\rho) = \frac{V_i \rho V_i^*}{\text{tr}(V_i \rho V_i^*)}$$

where the V_i are linear with $\sum_i V_i^* V_i = I$ and $p_i(\rho) = \text{tr}(V_i \rho V_i^*)$. Then Λ is written as

$$\Lambda(\rho) = \sum_i p_i F_i = \sum_i V_i \rho V_i^*$$

By simplicity we will assume that the quantum system considered can assume two states called 1 and 2.

We say that the pair $(\{X_n\}_{n \in \mathbb{N}}, \mu)$, $X_n : \Omega \rightarrow \{1, \dots, k\}$, is a **Quantum Stochastic Process**, QSP (homogeneous case), associated to the QIFS \mathcal{F} whenever μ is defined as

$$\mu(X_1 = x_1, \dots, X_n = x_n) := \text{tr}(V_{x_n} V_{x_{n-1}} \cdots V_{x_2} V_{x_1} \rho_0 V_{x_1}^* V_{x_2}^* \cdots V_{x_{n-1}}^* V_{x_n}^*) \quad (37)$$

where $\rho_0 \in \mathcal{M}_N$ is any density operator. The operator ρ_0 is a pre-measurement state, that is, we have a quantum system and we prepare ρ_0 as being its initial state (for a similar treatment to a sequence of measurements, see the definition of finite dimensional distributions in section 3).

So we can define for any r ,

$$\mu(X_r = x_r | X_{r-1} = x_{r-1}) = \frac{\text{tr}(V_{x_r} V_{x_{r-1}} \rho_0 V_{x_{r-1}}^* V_{x_r}^*)}{\text{tr}(V_{x_{r-1}} \rho_0 V_{x_{r-1}}^*)} \quad (38)$$

Definition 17. We say that a QSP is **Markov** if

$$\mu(X_n = x_n | X_1 = x_1, \dots, X_{n-1} = x_{n-1}) = \mu(X_n = x_n | X_{n-1} = x_{n-1}) \quad (39)$$

◇

Remark The condition $\sum_i V_i^* V_i = I$ is enough to show that the measure of a partition of cylinder sets equals 1. For instance, for two states 1 and 2, for $k = 2$ and writing

$$\mu(\overline{ij}) := \mu(X_1 = i, X_2 = j),$$

we have

$$\begin{aligned} & \mu(\overline{11}) + \mu(\overline{12}) + \mu(\overline{21}) + \mu(\overline{22}) \\ &= \text{tr}(V_1 V_1 \rho V_1^* V_1^*) + \text{tr}(V_2 V_1 \rho V_1^* V_2^*) + \text{tr}(V_1 V_2 \rho V_2^* V_1^*) + \text{tr}(V_2 V_2 \rho V_2^* V_2^*) \\ &= \text{tr}(V_1^* V_1 [V_1 \rho V_1^*]) + \text{tr}(V_2^* V_2 [V_1 \rho V_1^*]) + \text{tr}(V_1^* V_1 [V_2 \rho V_2^*]) + \text{tr}(V_2^* V_2 [V_2 \rho V_2^*]) \\ &= \text{tr}\left((V_1^* V_1 + V_2^* V_2) [V_1 \rho V_1^*]\right) + \text{tr}\left((V_1^* V_1 + V_2^* V_2) [V_2 \rho V_2^*]\right) \\ &= \text{tr}(V_1 \rho V_1^*) + \text{tr}(V_2 \rho V_2^*) = \text{tr}((V_1^* V_1 + V_2^* V_2) \rho) = 1 \end{aligned} \quad (40)$$

However, we note that there exist examples in which we can show that the measure of a partition of cylinder sets equals 1 even if we do not suppose that $\sum_i V_i^* V_i = I$. This happens, for instance, in the following construction involving stochastic matrices.

◇

Let us consider the particular case in which the operator $\rho_0 \in \mathcal{M}_N$, given in the definition of QSP is a fixed point for $\Lambda(\rho) = \sum_{i=1}^k V_i \rho V_i^*$ induced by the QIFS \mathcal{F} .

Suppose that V_1 and V_2 are defined by (31). Suppose that the matrix $P = (p_{ij})$ is column stochastic and that we have π such that $P\pi = \pi$. For instance we have

$$\mu(X_1 = 1, X_2 = 2) = \text{tr}(V_2 V_1 \rho_0 V_1^* V_2^*) = p_{21}(p_{11}\rho_{11} + p_{12}\rho_{22}) = p_{21}\rho_{11} \quad (41)$$

because with the choice of V_i we made, we have that the nonzero entries of ρ_0 correspond to the entries of π . So we can interpret p_{ij} as being

$$p_{ij} = \mu(X_2 = j | X_1 = i) \quad (42)$$

In a similar way,

$$\mu(X_1 = 2, X_2 = 1) = \text{tr}(V_1 V_2 \rho_0 V_2^* V_1^*) = p_{12}\rho_{22} \quad (43)$$

and

$$\mu(X_1 = 1, X_2 = 2, X_3 = 1) = \text{tr}(V_1 V_2 V_1 \rho_0 V_1^* V_2^* V_1^*) = p_{12}p_{21}\rho_{11} \quad (44)$$

Remark A simple calculation shows that with the V_i given by (31) we have that $\sum_i V_i^* V_i \neq I$. However, we still have that

$$\mu(\overline{11}) + \mu(\overline{12}) + \mu(\overline{21}) + \mu(\overline{22}) = 1$$

◇

To prove that the choice (31) reduces to the classic case for any sequence, we use the following lemma.

Lemma 1. *Suppose $N = 2$, $k = 2$. Then for every m , for V_i given by (31) and ρ_0 corresponding to the stationary vector π for P , we have that the product*

$$V_{x_m} V_{x_{m-1}} \cdots V_{x_1} \rho_0 V_{x_1}^* V_{x_2}^* \cdots V_{x_m}^* \quad (45)$$

has the form

$$\begin{pmatrix} * & 0 \\ 0 & 0 \end{pmatrix} \quad \text{ou} \quad \begin{pmatrix} 0 & 0 \\ 0 & * \end{pmatrix} \quad (46)$$

depending on whether $x_m = 1$ or $x_m = 2$, respectively.

Proof By induction. If $m = 1$ then

$$V_1 \rho_0 V_1^* = \begin{pmatrix} p_{11}\rho_{11} + p_{12}\rho_{22} & 0 \\ 0 & 0 \end{pmatrix} \quad (47)$$

and

$$V_2 \rho_0 V_2^* = \begin{pmatrix} 0 & 0 \\ 0 & p_{21}\rho_{11} + p_{22}\rho_{22} \end{pmatrix} \quad (48)$$

Suppose the lemma valid for m , we consider the product

$$V_{x_{m+1}} V_{x_m} \cdots V_{x_1} \rho_0 V_{x_1}^* V_{x_2}^* \cdots V_{x_m}^* V_{x_{m+1}}^* \quad (49)$$

Suppose $x_{m+1} = 1$. Then a simple calculation shows that

$$V_1 \begin{pmatrix} * & 0 \\ 0 & 0 \end{pmatrix} V_1^* \text{ e } V_1 \begin{pmatrix} 0 & 0 \\ 0 & * \end{pmatrix} V_1^* \quad (50)$$

has only one nonzero entry, namely the $(1, 1)$ -th entry. We proceed in a similar way for the case that $x_{m+1} = 2$, that is

$$V_2 \begin{pmatrix} * & 0 \\ 0 & 0 \end{pmatrix} V_2^* \text{ e } V_2 \begin{pmatrix} 0 & 0 \\ 0 & * \end{pmatrix} V_2^* \quad (51)$$

has only one nonzero entry, namely the $(2, 2)$ -th entry. □

Proposition 3. *If we set*

$$V_1 = \begin{pmatrix} \sqrt{p_{11}} & \sqrt{p_{12}} \\ 0 & 0 \end{pmatrix}, \quad V_2 = \begin{pmatrix} 0 & 0 \\ \sqrt{p_{21}} & \sqrt{p_{22}} \end{pmatrix}, \quad (52)$$

then

$$\mu(X_1 = x_1, X_2 = x_2, \dots, X_n = x_n) = p_{x_n x_{n-1}} p_{x_{n-1} x_{n-2}} \cdots p_{x_3 x_2} p_{x_2 x_1} \rho_{x_1 x_1} \quad (53)$$

where ρ_{ij} denotes the (i, j) -th entry of ρ_0 , eigenstate for $\Lambda(\rho) = \sum_i V_i \rho V_i^*$.

Proof We prove by induction. Suppose $n = 1$. Then

$$\mu(X_1 = 1) = \text{tr}(V_1 \rho_0 V_1^*) = p_{11} \rho_{11} + p_{12} \rho_{22} = \rho_{11}$$

$$\mu(X_1 = 2) = \text{tr}(V_2 \rho_0 V_2^*) = p_{21} \rho_{11} + p_{22} \rho_{22} = \rho_{22}$$

For the sake of clarity we also show the case $n = 2$. We have, after some routine calculations that

$$\mu(X_1 = 1, X_2 = 1) = \text{tr}(V_1 V_1 \rho_0 V_1^* V_1^*) = p_{11} \rho_{11} \quad (54)$$

$$\mu(X_1 = 1, X_2 = 2) = \text{tr}(V_2 V_1 \rho_0 V_1^* V_2^*) = p_{21} \rho_{11} \quad (55)$$

$$\mu(X_1 = 2, X_2 = 1) = \text{tr}(V_1 V_2 \rho_0 V_2^* V_1^*) = p_{12} \rho_{22} \quad (56)$$

$$\mu(X_1 = 2, X_2 = 2) = \text{tr}(V_2 V_2 \rho_0 V_2^* V_2^*) = p_{22} \rho_{22} \quad (57)$$

Now suppose the lemma holds for n , let us prove it for $n + 1$.

First suppose $x_{n+1} = 1$. Then

$$\begin{aligned} & \mu(X_1 = x_1, \dots, X_n = x_n, X_{n+1} = 1) \\ &= \text{tr}(V_1 V_{x_n} \cdots V_{x_2} V_{x_1} \rho_0 V_{x_1}^* V_{x_2}^* \cdots V_{x_{n-1}}^* V_{x_n}^* V_1^*) \end{aligned} \quad (58)$$

Using lemma 1, we have two cases. If $x_n = 1$ then

$$V_{x_n} \cdots V_{x_2} V_{x_1} \rho_0 V_{x_1}^* V_{x_2}^* \cdots V_{x_{n-1}}^* V_{x_n}^* = \begin{pmatrix} * & 0 \\ 0 & 0 \end{pmatrix}$$

and therefore

$$V_1 V_{x_n} \cdots V_{x_2} V_{x_1} \rho_0 V_{x_1}^* V_{x_2}^* \cdots V_{x_{n-1}}^* V_{x_n}^* V_1^* = V_1 \begin{pmatrix} * & 0 \\ 0 & 0 \end{pmatrix} V_1^* = \begin{pmatrix} *p_{11} & 0 \\ 0 & 0 \end{pmatrix}$$

and so by taking the trace we get

$$\begin{aligned} & tr(V_1 V_{x_n} \cdots V_{x_2} V_{x_1} \rho_0 V_{x_1}^* V_{x_2}^* \cdots V_{x_{n-1}}^* V_{x_n}^* V_1^*) \\ &= p_{11} p_{1x_{n-1}} p_{x_{n-1}x_{n-2}} \cdots p_{x_3x_2} p_{x_2x_1} \rho_{x_1x_1} \end{aligned} \quad (59)$$

In a similar way, if $x_n = 2$,

$$\begin{aligned} & V_{x_n} \cdots V_{x_2} V_{x_1} \rho_0 V_{x_1}^* V_{x_2}^* \cdots V_{x_{n-1}}^* V_{x_n}^* = \begin{pmatrix} 0 & 0 \\ 0 & * \end{pmatrix} \\ & V_1 V_{x_n} \cdots V_{x_2} V_{x_1} \rho_0 V_{x_1}^* V_{x_2}^* \cdots V_{x_{n-1}}^* V_{x_n}^* V_1^* = V_1 \begin{pmatrix} 0 & 0 \\ 0 & * \end{pmatrix} V_1^* = \begin{pmatrix} *p_{12} & 0 \\ 0 & 0 \end{pmatrix} \end{aligned}$$

and taking the trace gives

$$\begin{aligned} & tr(V_1 V_{x_n} \cdots V_{x_2} V_{x_1} \rho_0 V_{x_1}^* V_{x_2}^* \cdots V_{x_{n-1}}^* V_{x_n}^* V_1^*) \\ &= p_{12} p_{2x_{n-1}} p_{x_{n-1}x_{n-2}} \cdots p_{x_3x_2} p_{x_2x_1} \rho_{x_1x_1} \end{aligned} \quad (60)$$

Now we suppose $x_{n+1} = 2$, and we proceed in an analogous way.

$$\begin{aligned} & \mu(X_1 = x_1, \dots, X_n = x_n, X_{n+1} = 2) \\ &= tr(V_2 V_{x_n} \cdots V_{x_2} V_{x_1} \rho_0 V_{x_1}^* V_{x_2}^* \cdots V_{x_{n-1}}^* V_{x_n}^* V_2^*) \end{aligned} \quad (61)$$

By lemma 1, we have two cases. If $x_n = 1$ then

$$V_{x_n} \cdots V_{x_2} V_{x_1} \rho_0 V_{x_1}^* V_{x_2}^* \cdots V_{x_{n-1}}^* V_{x_n}^* = \begin{pmatrix} * & 0 \\ 0 & 0 \end{pmatrix}$$

therefore

$$V_2 V_{x_n} \cdots V_{x_2} V_{x_1} \rho_0 V_{x_1}^* V_{x_2}^* \cdots V_{x_{n-1}}^* V_{x_n}^* V_2^* = V_2 \begin{pmatrix} * & 0 \\ 0 & 0 \end{pmatrix} V_2^* = \begin{pmatrix} *p_{21} & 0 \\ 0 & 0 \end{pmatrix}$$

and taking the trace we get

$$\begin{aligned} & tr(V_2 V_{x_n} \cdots V_{x_2} V_{x_1} \rho_0 V_{x_1}^* V_{x_2}^* \cdots V_{x_{n-1}}^* V_{x_n}^* V_2^*) \\ &= p_{21} p_{1x_{n-1}} p_{x_{n-1}x_{n-2}} \cdots p_{x_3x_2} p_{x_2x_1} \rho_{x_1x_1} \end{aligned} \quad (62)$$

Analogously if $x_n = 2$

$$\begin{aligned} & V_{x_n} \cdots V_{x_2} V_{x_1} \rho_0 V_{x_1}^* V_{x_2}^* \cdots V_{x_{n-1}}^* V_{x_n}^* = \begin{pmatrix} 0 & 0 \\ 0 & * \end{pmatrix} \\ & V_2 V_{x_n} \cdots V_{x_2} V_{x_1} \rho_0 V_{x_1}^* V_{x_2}^* \cdots V_{x_{n-1}}^* V_{x_n}^* V_2^* = V_2 \begin{pmatrix} 0 & 0 \\ 0 & * \end{pmatrix} V_2^* = \begin{pmatrix} *p_{22} & 0 \\ 0 & 0 \end{pmatrix} \end{aligned}$$

and taking the trace

$$\begin{aligned} & \text{tr}(V_2 V_{x_n} \cdots V_{x_2} V_{x_1} \rho_0 V_{x_1}^* V_{x_2}^* \cdots V_{x_{n-1}}^* V_{x_n}^* V_2^*) \\ &= p_{22} p_{2x_{n-1}} p_{x_{n-1}x_{n-2}} \cdots p_{x_3x_2} p_{x_2x_1} \rho_{x_1x_1} \end{aligned} \quad (63)$$

□

Corollary 1. *The quantum stochastic process induced by*

$$V_1 = \begin{pmatrix} \sqrt{p_{11}} & \sqrt{p_{12}} \\ 0 & 0 \end{pmatrix}, \quad V_2 = \begin{pmatrix} 0 & 0 \\ \sqrt{p_{21}} & \sqrt{p_{22}} \end{pmatrix}, \quad (64)$$

is Markov.

Proof By the proposition, we have that the measure μ reduces to the Markov measure for matrices.

□

Lemma 2. *For V_i linear maps and ρ_0 fixed point for $\Lambda = \sum_i V_i \rho V_i^*$, we have for any m, n ,*

$$\begin{aligned} & \mu(X_1 = x_1, X_2 = x_2, \dots, X_n = x_n) = \\ & \mu(X_m = x_1, X_{m+1} = x_2, \dots, X_{m+n} = x_n) \end{aligned}$$

Proof We prove the lemma for the case in which we have two possible states 1 and 2. We have

$$\begin{aligned} & \mu(X_m = x_1, X_{m+1} = x_2, \dots, X_{m+n} = x_n) \\ &= \sum_{i_1, \dots, i_{m-1}} \mu(X_1 = i_1, X_2 = i_2, \dots, X_{m-1} = i_{m-1}, X_m = x_1, \dots, X_{m+n} = x_n) \\ &= \sum_{i_2, \dots, i_{m-1}} \text{tr}(V_{x_n} \cdots V_{x_1} V_{i_{m-1}} \cdots V_{i_2} V_1 \rho_0 V_1^* V_{i_2}^* \cdots) \\ & \quad + \text{tr}(V_{x_n} \cdots V_{x_1} V_{i_{m-1}} \cdots V_{i_2} V_2 \rho_0 V_2^* V_{i_2}^* \cdots) \\ &= \sum_{i_2, \dots, i_{m-1}} \text{tr}(V_{x_n} \cdots V_{x_1} V_{i_{m-1}} \cdots V_{i_2} \rho_0 V_{i_2}^* V_{i_3}^* \cdots V_{i_{m-1}}^* V_{x_1}^* \cdots V_{x_n}^*) \end{aligned}$$

Repeating the procedure above for i_2, i_3 , etc. we get

$$\mu(X_m = x_1, X_{m+1} = x_2, \dots, X_{m+n} = x_n) = \text{tr}(V_{x_n} \cdots V_{x_1} \rho_0 V_{x_1}^* \cdots V_{x_n}^*)$$

This concludes the proof.

□

Example 10. Let us make an inspection with respect to the Chapman-Kolmogorov equation, that is, we would like to know if the equality

$$\mu_{ij}(m+n) = \sum_k \mu_{ik}(m)\mu_{kj}(n) \quad (65)$$

holds, where

$$\mu_{ij}(n) = \mu(X_{m+n} = j | X_m = i)$$

Take for instance, $m = n = i = j = 1$. Then

$$\begin{aligned} \sum_k \mu_{ik}(m)\mu_{kj}(n) &= \mu_{11}(1)\mu_{11}(1) + \mu_{12}(1)\mu_{21}(1) \\ &= \frac{\text{tr}(V_1 V_1 \rho V_1^* V_1^*)^2}{\text{tr}(V_1 \rho V_1^*)^2} + \frac{\text{tr}(V_2 V_1 \rho V_1^* V_2^*)}{\text{tr}(V_1 \rho V_1^*)} \frac{\text{tr}(V_1 V_2 \rho V_2^* V_1^*)}{\text{tr}(V_2 \rho V_2^*)} \end{aligned} \quad (66)$$

and

$$\begin{aligned} \mu_{ij}(m+n) &= \mu_{11}(2) = \mu(X_3 = 1 | X_1 = 1) \\ &= \frac{\text{tr}(V_1 V_1 V_1 \rho V_1^* V_1^* V_1^*)}{\text{tr}(V_1 \rho V_1^*)} + \frac{\text{tr}(V_1 V_2 V_1 \rho V_1^* V_2^* V_1^*)}{\text{tr}(V_1 \rho V_1^*)} \end{aligned} \quad (67)$$

Now let V_1, V_2 be given by (52), then we obtain classic calculations, so the Chapman-Kolmogorov equation holds. Now take

$$V_1 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad V_2 = \begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix} \quad (68)$$

then we get, from (66) and (67):

$$\frac{\text{tr}(V_1 V_1 \rho V_1^* V_1^*)^2}{\text{tr}(V_1 \rho V_1^*)^2} + \frac{\text{tr}(V_2 V_1 \rho V_1^* V_2^*)}{\text{tr}(V_1 \rho V_1^*)} \frac{\text{tr}(V_1 V_2 \rho V_2^* V_1^*)}{\text{tr}(V_2 \rho V_2^*)} = 1 + \frac{\rho_{11}}{\rho_{11} + 4\rho_{22}} \quad (69)$$

and

$$\frac{\text{tr}(V_1 V_1 V_1 \rho V_1^* V_1^* V_1^*)}{\text{tr}(V_1 \rho V_1^*)} + \frac{\text{tr}(V_1 V_2 V_1 \rho V_1^* V_2^* V_1^*)}{\text{tr}(V_1 \rho V_1^*)} = 1 + 1 = 2 \quad (70)$$

Then in this case we have that the Chapman-Kolmogorov equation holds if and only if $\rho_{22} = 0$ that is, if $\rho_{11} = 1$. Also, we note that $\sum_i V_i^* V_i \neq I$. To conclude this example, we take V_1 and V_2 with $\sum_i V_i^* V_i = I$, namely,

$$V_1 = \begin{pmatrix} \frac{1}{\sqrt{3}} & 0 \\ 0 & 0 \end{pmatrix}, \quad V_2 = \begin{pmatrix} \sqrt{\frac{2}{3}} & 0 \\ 0 & 1 \end{pmatrix} \quad (71)$$

Take for instance $\rho_0 = \frac{1}{4}|1\rangle\langle 1| + \frac{3}{4}|2\rangle\langle 2|$, a fixed point for the associated Λ . A simple calculation shows that (66) and (67) are different. Therefore our calculation shows that the Chapman-Kolmogorov equation does not hold in general (for our setting).

◇

We would like to obtain a nonhomogeneous version for the measure we defined in (37) in the homogeneous case, i.e., we are looking for a measure induced by a nonhomogeneous QIFS. Let W_i , $i = 1, \dots, k$ be linear operators such that $\sum_i W_i^* W_i = I$. Let $\rho_0 \in \mathcal{M}_N$. Define

$$\begin{aligned} \mu(X_1 = x_1, \dots, X_n = x_n) &:= \\ &= \text{tr}(W_{x_1} \rho_0 W_{x_1}^*) \frac{\text{tr}(W_{x_2} V_{x_1} \rho_0 V_{x_1}^* W_{x_2}^*)}{\text{tr}(V_{x_1} \rho_0 V_{x_1}^*)} \frac{\text{tr}(W_{x_3} V_{x_2} V_{x_1} \rho_0 V_{x_1}^* V_{x_2}^* W_{x_3}^*)}{\text{tr}(V_{x_2} V_{x_1} \rho_0 V_{x_1}^* V_{x_2}^*)} \times \dots \\ &\dots \times \frac{\text{tr}(W_{x_{n-1}} V_{x_{n-2}} \dots V_{x_1} \rho_0 V_{x_1}^* \dots V_{x_{n-2}}^* W_{x_{n-1}}^*)}{\text{tr}(V_{x_{n-2}} \dots V_{x_1} \rho_0 V_{x_1}^* \dots V_{x_{n-2}}^*)} \times \\ &\times \frac{\text{tr}(W_{x_n} V_{x_{n-1}} \dots V_{x_1} \rho_0 V_{x_1}^* \dots V_{x_{n-1}}^* W_{x_n}^*)}{\text{tr}(V_{x_{n-1}} \dots V_{x_1} \rho_0 V_{x_1}^* \dots V_{x_{n-1}}^*)} \end{aligned} \quad (72)$$

that is,

$$\mu(X_1 = x_1, \dots, X_n = x_n) := \text{tr}(W_{x_1} \rho_0 W_{x_1}^*) \prod_{i=2}^n \frac{\text{tr}(W_{x_i} V_{x_{i-1}} \dots V_{x_1} \rho_0 V_{x_1}^* \dots V_{x_{i-1}}^* W_{x_i}^*)}{\text{tr}(V_{x_{i-1}} V_{x_{i-2}} \dots V_{x_1} \rho_0 V_{x_1}^* \dots V_{x_{i-2}}^* V_{x_{i-1}}^*)} \quad (73)$$

Remark A calculation shows that if we suppose $\sum_i W_i^* W_i = I$, then

$$\sum_{i_1, \dots, i_n} \mu(\overline{i_1 \dots i_n}) = 1$$

Besides, if we suppose that $W_i = V_i$ for all i , then we recover the measure definition for homogeneous QSP.

◇

Consider a QIFS $\mathcal{F} = \{\mathcal{M}_N, F_i, p_i\}_{i=1, \dots, k}$, where

$$F_i(\rho) = \frac{V_i \rho V_i^*}{\text{tr}(V_i \rho V_i^*)}$$

where the V_i are linear and $p_i(\rho) = \text{tr}(W_i \rho W_i^*)$, com $\sum_i W_i^* W_i = I$.

Definition 18. We say that the pair $(\{X_n\}_{n \in \mathbb{N}}, \mu)$, $X_n : \Omega \rightarrow \{1, \dots, k\}$, is a **Quantum Stochastic Process** associated to the **nonhomogeneous QIFS** \mathcal{F} if μ is defined by (73), where $\rho_0 \in \mathcal{M}_N$ is any density operator.

Remark In the definition above we can, of course, consider the particular case in which ρ_0 is a fixed point for

$$\Lambda(\rho) = \sum_{i=1}^k \text{tr}(W_i \rho W_i^*) \frac{V_i \rho V_i^*}{\text{tr}(V_i \rho V_i^*)},$$

induced by the QIFS \mathcal{F} .

◇

Recall that by lemma 2, a homogeneous QSP is always stationary. This is no longer true in general for nonhomogeneous QSP.

Example 11. Let $\{X_n\}_{n \in \mathbb{N}}$ be a QSP induced by a nonhomogeneous QIFS. We would like to know whether

$$\mu(X_1 = 1, X_2 = 2) = \mu(X_2 = 1, X_3 = 2) \quad (74)$$

By definition we have:

$$\mu(X_1 = 1, X_2 = 2) = \text{tr}(W_1 \rho_0 W_1^*) \frac{\text{tr}(W_2 V_1 \rho_0 V_1^* W_2^*)}{\text{tr}(V_1 \rho_0 V_1^*)} \quad (75)$$

And also

$$\begin{aligned} \mu(X_2 = 1, X_3 = 2) &= \mu(X_1 = 1, X_2 = 1, X_3 = 2) + \mu(X_1 = 2, X_2 = 1, X_3 = 2) \\ &= \text{tr}(W_1 \rho_0 W_1^*) \frac{\text{tr}(W_1 V_1 \rho_0 V_1^* W_1^*)}{\text{tr}(V_1 \rho_0 V_1^*)} \frac{\text{tr}(W_2 V_1 V_1 \rho_0 V_1^* V_1^* W_2^*)}{\text{tr}(V_1 V_1 \rho_0 V_1^* V_1^*)} \\ &\quad + \text{tr}(W_2 \rho_0 W_2^*) \frac{\text{tr}(W_1 V_2 \rho_0 V_2^* W_1^*)}{\text{tr}(V_2 \rho_0 V_2^*)} \frac{\text{tr}(W_2 V_1 V_2 \rho_0 V_2^* V_1^* W_2^*)}{\text{tr}(V_1 V_2 \rho_0 V_2^* V_1^*)} \end{aligned} \quad (76)$$

$$\begin{aligned} &= \text{tr} \left[W_2 V_1 \left[\text{tr}(W_1 \rho_0 W_1^*) \frac{V_1 \rho_0 V_1^*}{\text{tr}(V_1 \rho_0 V_1^*)} \left(\frac{\text{tr}(W_1 V_1 \rho_0 V_1^* W_1^*)}{\text{tr}(V_1 V_1 \rho_0 V_1^* V_1^*)} \right) + \right. \right. \\ &\quad \left. \left. + \text{tr}(W_2 \rho_0 W_2^*) \frac{V_2 \rho_0 V_2^*}{\text{tr}(V_2 \rho_0 V_2^*)} \left(\frac{\text{tr}(W_1 V_2 \rho_0 V_2^* W_1^*)}{\text{tr}(V_1 V_2 \rho_0 V_2^* V_1^*)} \right) \right] V_1^* W_2^* \right] \end{aligned} \quad (77)$$

Note that in the homogeneous case we have that both fractions in parenthesis on equation (77) are equal to 1, so if ρ_0 is a fixed point for Λ , then we have stationarity, a fact we have already proved. But in the nonhomogeneous case, the terms in parenthesis are not equal to 1 in general.

◇

8. A definition of entropy for QIFS

We will present a notion of entropy for “invariant” (or “stationary”) measures with support on density matrices. This definition is obtained by adapting the reasoning described in [5], [12] and [13] to the present situation. The main idea is to define this concept via the Ruelle operator and to avoid the use of partitions.

Denote by p an arbitrary choice of mappings $p_i : \mathcal{M}_N \rightarrow \mathbb{R}$, $i = 1, \dots, k$ for a certain k . Let

$$m_b(\mathcal{M}_N) := \{f : \mathcal{M}_N \rightarrow \mathbb{R} : f \text{ is measurable and bounded}\}$$

Let $\mathcal{U}_p : m_b(\mathcal{M}_N) \rightarrow m_b(\mathcal{M}_N)$,

$$(\mathcal{U}_p f)(\rho) := \sum_{i=1}^k p_i(\rho) f(F_i(\rho))$$

Let us consider all possible choices of mappings $p_i : \mathcal{M}_N \rightarrow \mathbb{R}$ which satisfy

$$\mathcal{U}_p 1 = 1 \quad (78)$$

Each p determines an operator \mathcal{U}_p . The set of all possible p that satisfy (78) will be denoted by P .

Let $(\mathcal{M}_N, F_i, p_i)_{i=1, \dots, k}$ be a QIFS. An example of Markov operator for measures is the one we defined before, given by $\mathcal{V}_p : M^1(\mathcal{M}_N) \rightarrow M^1(\mathcal{M}_N)$,

$$(\mathcal{V}_p \nu)(B) = \sum_{i=1}^k \int_{F_i^{-1}(B)} p_i d\nu,$$

which we will call **the Markov operator Markov induced by the p_i** . That is, we will consider all \mathcal{V}_p with $p \in P$. We say that ν is **invariant** for the F_i if for some $p \in P$ we have that $\mathcal{V}_p \nu = \nu$.

Let \mathcal{M}_F be the set of all invariant measures for a fixed choice of the dynamics F_i , $i = 1, \dots, k$. For such measures $\nu \in \mathcal{M}_F$, and based on [5], [12] and [13], define

$$h_0(\nu) := \inf_{f \in \mathbb{B}^+} \int \log \left(\sum_{i=1}^k \frac{f \circ F_i}{f} \right) d\nu$$

Above, \mathbb{B}^+ denotes the bounded, positive, borelean functions on \mathcal{M}_N .

Proposition 4. *For $\nu \in \mathcal{M}_F$, we have that $0 \leq h_0(\nu) \leq \log k$.*

In order to prove this proposition, we need the following lemma.

Lemma 3. [13] *Let $\beta \geq 1 + \alpha$ and numbers $a_i \in [1 + \alpha, \beta]$, $i = 1, \dots, k$. Then there exists $\epsilon \geq 1$ such that*

$$\log \left(\epsilon \sum_{i=1}^k a_i \right) \geq \sum_{i=1}^k \log (\epsilon a_i).$$

The proof of this lemma follows by choosing

$$\epsilon = \exp\left(\frac{1}{k} \frac{\log \sum_{i=1}^k a_i}{\sum_{i=1}^k \log a_i}\right)$$

Lemma 4. *If $f \in \mathbb{B}^+$ and $\nu \in \mathcal{M}_F$ then*

$$\sum_{i=1}^k \int f \circ F_i d\nu \geq \int f d\nu$$

Proof First suppose that $f = 1_B$, where B is a measurable set. We have that

$$\begin{aligned} \sum_{i=1}^k \int 1_B \circ F_i d\nu &\geq \sum_{i=1}^k \int p_i(x) 1_B(F_i(x)) d\nu(x) = \sum_{i=1}^k \int_{F_i^{-1}(B)} p_i(x) d\nu(x) \\ &= \mathcal{V}_p(\nu)(B) = \nu(B) = \int 1_B d\nu \end{aligned}$$

Then, assume that $f = \sum_{j=1}^l b_j 1_{B_j}$, i.e., a simple function. Then

$$\begin{aligned} \sum_{i=1}^k \int \sum_{j=1}^l b_j 1_{B_j} \circ F_i d\nu &= \sum_{j=1}^l b_j \sum_{i=1}^k \int 1_{B_j} \circ F_i d\nu \\ &\geq \sum_{j=1}^l b_j \sum_{i=1}^k \int p_i(x) 1_{B_j}(F_i(x)) d\nu = \sum_{j=1}^l b_j \mathcal{V}_p(\nu)(B_j) \\ &= \sum_{j=1}^l b_j \nu(B_j) = \int f d\nu \end{aligned}$$

Now let $f = \lim_n f_n$, a limit of a sequence of simple functions. Note that we suppose $f \in \mathbb{B}^+$, so f is bounded, and since ν is a probability measure on \mathcal{M}_N , it follows that f is integrable. By the bounded convergence theorem, we have that

$$\begin{aligned} \sum_{i=1}^k \int f \circ F_i d\nu &= \sum_{i=1}^k \int \lim_n f_n \circ F_i d\nu = \lim_n \sum_{i=1}^k \int f_n \circ F_i d\nu \\ &\geq \lim_n \int f_n d\nu = \int \lim_n f_n d\nu = \int f d\nu \end{aligned}$$

□

The following proof is an adaptation of results seen in [13].

Proof of proposition 4 Let us restrict the proof for the case in which we have a QIFS $(\mathcal{M}_N, F_i, p_i)_{i=1, \dots, k}$, where $F_i(\rho) = V_i \rho V_i^*$, with linear V_i .

First note that if $f \equiv 1$, we have $\int \log(\sum_{i=1}^k 1) d\nu = \log k$, so $h_0(\nu) \leq \log k$.

Let $I = \int \log(\sum_{i=1}^k \frac{f \circ F_i}{f}) d\nu$ and suppose, without loss of generality, that $1 + \alpha \leq f \leq \beta$ (note that this integral is invariant by the projective mapping $f \rightarrow \lambda f$). Then

$$I = \int \log\left(\sum_{i=1}^k \frac{\epsilon f \circ F_i}{\epsilon f}\right) d\nu = \int \log\left(\sum_{i=1}^k \epsilon f \circ F_i\right) d\nu - \int \log(\epsilon f) d\nu \quad (79)$$

Define

$$a_i = f \circ F_i(\rho)$$

Then

$$\epsilon(\rho) = \exp\left(\frac{1}{k} \frac{\log \sum_{i=1}^k f \circ F_i}{\log f \circ F_i}\right) \geq \epsilon_0 \geq 1,$$

by the compactness of \mathcal{M}_N . With such choice we obtain, by lemma (3),

$$\log\left(\epsilon_0 \sum_{i=1}^k f \circ F_i\right) \geq \sum_{i=1}^k \log(\epsilon_0 f \circ F_i) \quad (80)$$

Apply (80) on (79), then

$$I \geq \sum_{i=1}^k \int \log(\epsilon_0 f \circ F_i) d\nu - \int \log(\epsilon_0 f) d\nu$$

Then by lemma (4) applied on the function $\log(\epsilon f)$ (note that we have $\log(\epsilon_0 f) \in \mathbb{B}^+$ because $\epsilon_0 \geq 1$), we get

$$I \geq \int \log(\epsilon f) d\nu - \int \log(\epsilon f) d\nu = 0$$

□

The computation in the next example shows that the concept of entropy described here is different from the one presented in [1] [2].

Example 12. We will consider an example of a probability η such that $\mathcal{V}(\eta) = \eta$ and we will compute the entropy of η .

Suppose a QIFS, such that

$$p_i(\rho) = \text{tr}(W_i \rho W_i^*), \quad \sum_i W_i^* W_i = I, \quad F_i(\rho) = \frac{V_i \rho V_i^*}{\text{tr}(V_i \rho V_i^*)}$$

for $i = 1, \dots, k$. Denote $m_b(\mathcal{M}_N)$ the space of bounded and measurable functions in \mathcal{M}_N .

Consider $\Lambda : \mathcal{M}_N \rightarrow \mathcal{M}_N$,

$$\Lambda(\rho) = \sum_i p_i(\rho) F_i(\rho) = \sum_i \text{tr}(W_i \rho W_i^*) \frac{V_i \rho V_i^*}{\text{tr}(V_i \rho V_i^*)}$$

Suppose there exists a density matrix ρ which Λ -invariant. As we know, such state is the barycenter of μ which is \mathcal{V} -invariant [1].

Suppose $\mathcal{V}\mu = \mu$, then we can write

$$\begin{aligned} \int f d\mu &= \int f d\mathcal{V}\mu = \sum_{i=1}^k \int p_i(\rho) f(F_i(\rho)) d\mu(\rho) = \\ &= \sum_i \int p_i(\rho) f\left(\frac{V_i \rho V_i^*}{\text{tr}(V_i \rho V_i^*)}\right) d\mu = \sum_i \int \text{tr}(W_i \rho W_i^*) f\left(\frac{V_i \rho V_i^*}{\text{tr}(V_i \rho V_i^*)}\right) d\mu \end{aligned}$$

Therefore, for any $f \in m_b(\mathcal{M}_N)$, we got the condition

$$\int f d\mu = \sum_i \int \text{tr}(W_i \rho W_i^*) f\left(\frac{V_i \rho V_i^*}{\text{tr}(V_i \rho V_i^*)}\right) d\mu \quad (81)$$

Let us consider a particular example where $N = 2$, $k = 4$, and

$$\begin{aligned} V_1 &= \begin{pmatrix} \sqrt{p_{11}} & 0 \\ 0 & 0 \end{pmatrix}, \quad V_2 = \begin{pmatrix} 0 & \sqrt{p_{12}} \\ 0 & 0 \end{pmatrix}, \\ V_3 &= \begin{pmatrix} 0 & 0 \\ \sqrt{p_{21}} & 0 \end{pmatrix}, \quad V_4 = \begin{pmatrix} 0 & 0 \\ 0 & \sqrt{p_{22}} \end{pmatrix}, \end{aligned}$$

in such way that the p_{ij} are the entries of a column stochastic matrix P . Let $\pi = (\pi_1, \pi_2)$ be a vector such that $P\pi = \pi$. A simple calculation shows that for ρ , the density matrix such that has entries ρ_{ij} , we have

$$V_1 \rho V_1^* = \begin{pmatrix} p_{11} \rho_{11} & 0 \\ 0 & 0 \end{pmatrix}, \quad V_2 \rho V_2^* = \begin{pmatrix} p_{12} \rho_{22} & 0 \\ 0 & 0 \end{pmatrix} \quad (82)$$

$$V_3 \rho V_3^* = \begin{pmatrix} 0 & 0 \\ 0 & p_{21} \rho_{11} \end{pmatrix}, \quad V_4 \rho V_4^* = \begin{pmatrix} 0 & 0 \\ 0 & p_{22} \rho_{22} \end{pmatrix}, \quad (83)$$

and therefore

$$\frac{V_1 \rho V_1^*}{\text{tr}(V_1 \rho V_1^*)} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad \frac{V_2 \rho V_2^*}{\text{tr}(V_2 \rho V_2^*)} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \quad (84)$$

$$\frac{V_3 \rho V_3^*}{\text{tr}(V_3 \rho V_3^*)} = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \quad \frac{V_4 \rho V_4^*}{\text{tr}(V_4 \rho V_4^*)} = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \quad (85)$$

that is, the above values do not depend on ρ .

Define

$$\rho_x = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad \rho_y = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \quad (86)$$

and

$$\eta = \pi_1 \delta_{\rho_x} + \pi_2 \delta_{\rho_y} \quad (87)$$

Note that the barycenter of η is

$$\rho_\eta = \pi_1 \rho_x + \pi_2 \rho_y = \pi_1 \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + \pi_2 \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} \pi_1 & 0 \\ 0 & \pi_2 \end{pmatrix}$$

One can show directly that $\mathcal{V}(\eta) = \eta$ (see [1]). Define

$$\rho_1 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad \rho_2 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \quad (88)$$

and also

$$\eta = \pi_1 \delta_{\rho_1} + \pi_2 \delta_{\rho_2} \quad (89)$$

Note that the barycenter of η is

$$\rho_\eta = \pi_1 \rho_1 + \pi_2 \rho_2 = \pi_1 \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} + \pi_2 \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} \pi_1 & 0 \\ 0 & \pi_2 \end{pmatrix}$$

From this it will also follow that $\mathcal{V}\eta = \eta$ [1]. We will show that the entropy of such η is $\log(2) - \pi_1 \log(\pi_1) - \pi_2 \log(\pi_2)$. Remember that

$$\begin{aligned} & \int \log \left(\sum_i \frac{f \circ F_i}{f} \right) d\mu \\ &= \int \log \left(\sum_i f \left(\frac{V_i \rho V_i^*}{\text{tr}(V_i \rho V_i^*)} \right) \right) d\mu - \sum_i \int \text{tr}(W_i \rho W_i^*) \log f \left(\frac{V_i \rho V_i^*}{\text{tr}(V_i \rho V_i^*)} \right) d\mu. \end{aligned} \quad (90)$$

For such choice of V_i take

$$c_i = f \left(\frac{V_i \rho V_i^*}{\text{tr}(V_i \rho V_i^*)} \right), \quad i = 1, \dots, 4 \quad (91)$$

Note that

$$c_1 = c_2, \quad c_3 = c_4 \quad (92)$$

Then we can write

$$\int \log \left(\sum_i \frac{f \circ F_i}{f} \right) d\eta = \int \log \left(\sum_i c_i \right) d\eta - \sum_i \int \operatorname{tr}(W_i \rho W_i^*) \log c_i d\eta. \quad (93)$$

Therefore

$$\begin{aligned} & \int \log \left(\sum_i \frac{f \circ F_i}{f} \right) d\eta = \pi_1 \log \left(\sum_i c_i \right) + \pi_2 \log \left(\sum_i c_i \right) \\ & \quad - \sum_i \left[\operatorname{tr}(V_i \rho_1 V_i^*) \pi_1 \log(c_i) + \operatorname{tr}(V_i \rho_2 V_i^*) \pi_2 \log(c_i) \right] \\ & \quad = \pi_1 \log(2c_1 + 2c_3) + \pi_2 \log(2c_1 + 2c_3) \\ & \quad - \sum_i \left[\operatorname{tr}(V_i \rho_1 V_i^*) \pi_1 \log(c_i) + \operatorname{tr}(V_i \rho_2 V_i^*) \pi_2 \log(c_i) \right] \\ & = \log(2(c_1 + c_3)) - \sum_i \left[\operatorname{tr}(V_i \rho_1 V_i^*) \pi_1 \log(c_i) + \operatorname{tr}(V_i \rho_2 V_i^*) \pi_2 \log(c_i) \right] \\ & \quad = \log(2(c_1 + c_3)) \\ & \quad - \left[\pi_1 \left(p_{11}^2 \log(c_1) + p_{12} p_{21} \log(c_2) + p_{21} p_{11} \log(c_3) + p_{22} p_{21} \log(c_4) \right) \right. \\ & \quad \left. + \pi_2 \left(p_{11} p_{12} \log(c_1) + p_{12} p_{22} \log(c_2) + p_{21} p_{12} \log(c_3) + p_{22}^2 \log(c_4) \right) \right] \\ & \quad = \log(2(c_1 + c_3)) \\ & \quad - \left[p_{11} \log(c_1) (\pi_1 p_{11} + \pi_2 p_{12}) + p_{12} \log(c_2) (\pi_1 p_{21} + \pi_2 p_{22}) \right. \\ & \quad \left. + p_{21} \log(c_3) (\pi_1 p_{11} + \pi_2 p_{12}) + p_{22} \log(c_4) (\pi_1 p_{21} + \pi_2 p_{22}) \right] \\ & \quad = \log(2(c_1 + c_3)) \\ & \quad - \left[\pi_1 p_{11} \log(c_1) + \pi_2 p_{12} \log(c_2) + \pi_1 p_{21} \log(c_3) + \pi_2 p_{22} \log(c_4) \right] \\ & \quad = \log(2(c_1 + c_3)) - (\pi_1 \log(c_1) + \pi_2 \log(c_3)) \end{aligned}$$

Finally,

$$\int \log \left(\sum_i \frac{f \circ F_i}{f} \right) d\eta = \log(2(c_1 + c_3)) - (\pi_1 \log(c_1) + \pi_2 \log(c_3)). \quad (94)$$

Now we will use Lagrange multipliers. Define $b : \mathbb{R}_+^2 \rightarrow \mathbb{R}$, where \mathbb{R}_+^2 is the set of positive coordinates, by

$$b(x, y) = \log(2(x + y)) - (\pi_1 \log(x) + \pi_2 \log(y))$$

We impose the restriction

$$x + y = a$$

for fixed $a > 0$. We will get below the critical point of b under such restriction. After that we consider a general $a > 0$.

Define

$$g(x, y) = x + y - a$$

and

$$\Gamma(x, y, \lambda) = b + \lambda g$$

Then, $\nabla \Gamma = 0$ implies

$$\frac{1}{x+y} - \frac{\pi_1}{x} + \lambda = 0 \quad (95)$$

$$\frac{1}{x+y} - \frac{\pi_2}{y} + \lambda = 0 \quad (96)$$

$$x + y = a \quad (97)$$

from which follows

$$x = \pi_1 a, \quad y = \pi_2 a. \quad (98)$$

Therefore,

$$c_1 = c_2 = \pi_1 a, \quad c_3 = c_4 = \pi_2 a \quad (99)$$

From (94) we get

$$\begin{aligned} \int \log \left(\sum_i \frac{f \circ F_i}{f} \right) d\eta &= \log(2(\pi_1 a + \pi_2 a)) - (\pi_1 \log(\pi_1 a) + \pi_2 \log(\pi_2 a)) \\ &= \log(2a) - \pi_1 \log(\pi_1 a) - \pi_2 \log(\pi_2 a) \\ &= \log(2) + \log(a) - \pi_1 \log(\pi_1) - \pi_1 \log(a) - \pi_2 \log(\pi_2) - \pi_2 \log(a) \\ &= \log(2) - \pi_1 \log(\pi_1) - \pi_2 \log(\pi_2) \end{aligned} \quad (100)$$

This value of entropy is different from the value computed in the same example of QIFS in [1], [2] which is $-\sum_{i,j} \pi_i p_{ji} \log p_{ji}$ (Example 7 in section 11 [1]).

◇

Given the expression

$$h_0(\nu) := \inf_{f \in \mathbb{B}^+} \int \log \left(\sum_{i=1}^k \frac{f \circ F_i}{f} \right) d\nu,$$

for a fixed probability ν , which is invariant by the shift acting on the space Ω , a natural question is to identify the f which realizes the infimum above.

We will describe below the analysis of the classical case (in the sense of Stochastic Processes, and not QSP). Our purpose is to explain why

the definition presented above is a natural generalization of the setting for Markov Processes. In the case the probability ν comes from a Markov Process this will be now derived.

Let $\Omega = I_m^{\mathbb{N}}$, where $I_m = \{1, \dots, m\}$, and let $\mathcal{C} = \{C_\iota : \iota \in \cup_{n \in \mathbb{N}} I_m^n\}$ the collection of cylinder sets in Ω , where

$$C_\iota := \{\omega \in I_m^{\mathbb{N}} : w(j) = i_j, j = 1, \dots, r, \iota = (i_1, \dots, i_r) \in I_m^r\}$$

and denote by $\sigma(\mathcal{C})$ the σ -algebra generated by the cylinders in Ω .

Let (P, π) be a Markov chain, so that $P = (p_{ij})$ is a matrix of order n , with $p_{ij} \geq 0$, $\sum_j p_{ij} = 1$ (row stochastic), and $\pi = (\pi_1, \dots, \pi_n)$ is the left eigenvector with eigenvalue 1. So $\pi P = \pi$, that is, $\sum_i \pi_i p_{ij} = \pi_j$.

Associated to the matrix P we have the following measure.

Definition 19. *The Markov measure (associated to the chain (P, π)) of a cylinder is defined as*

$$\nu(C_\iota) := \pi_{i_1} p_{i_1 i_2} p_{i_2 i_3} \cdots p_{i_{r-1} i_r} \quad (101)$$

◇

We are interested in the following problem: find the infimum f in

$$h_0(\nu) := \inf_{f \in \mathbb{B}^+} \int \log\left(\sum_{i=1}^k \frac{f \circ F_i}{f}\right) d\nu \quad (102)$$

for such ν defined above.

◇

We use the notation \overline{ij} to denote the cylinder set in $I_m^{\mathbb{N}}$ which consists of the set of sequences (w_1, w_2, \dots) such that $w_1 = i$ and $w_2 = j$. Denote by $1_{\overline{ij}}$ the indicator function of \overline{ij} . To simplify, suppose $m = 2$ so the alphabet considered contains only two symbols, denoted by 1 and 2. Define the following function $f : I_2^{\mathbb{N}} \rightarrow \mathbb{R}^+$,

$$f(x) = \sum_{i,j=1}^2 a_{ij} 1_{\overline{ij}}(x) \quad (103)$$

where $a_{ij} \in \mathbb{R}^+$. That is, f is a simple function, constant on \overline{ij} . In this form, $\log f = \sum_{i,j} \log a_{ij} 1_{\overline{ij}}$.

Let us suppose that $F_i : I_m^{\mathbb{N}} \rightarrow I_m^{\mathbb{N}}$ is the mapping $F_i(w_1, w_2, \dots) = (i, w_1, w_2, \dots)$. If ν is a Markov measure, we have

$$\int_{I_2^{\mathbb{N}}} \log f d\nu = \int_{I_2^{\mathbb{N}}} \sum_{i,j=1}^2 \log(a_{ij}) 1_{\bar{ij}} d\nu = \sum_{i,j=1}^2 \log(a_{ij}) \nu(\bar{ij}) = \sum_{i,j=1}^2 \pi_i p_{ij} \log a_{ij} \quad (104)$$

Also, we have for $w = (i, j, \dots)$,

$$f \circ F_l(w) = \sum_{i,j} a_{ij} 1_{\bar{ij}}(F_l(w)) = a_{li} \quad (105)$$

To see that, note that by the expression above we have a sum of terms such that $(i, j) = (l, i)$, therefore $a_{ij} = a_{li}$.

Then

$$\begin{aligned} \int \log\left(\sum_{i=1}^2 \frac{f \circ F_i}{f}\right) d\nu &= \int \log\left(\sum_{l=1}^2 f \circ F_l\right) d\nu - \int \log f d\nu \\ &= \int \log\left(\sum_{l=1}^2 f \circ F_l\right) d\nu - \sum_{i,j=1}^2 \pi_i p_{ij} \log a_{ij} \end{aligned} \quad (106)$$

Note that for any $w \in I_m^{\mathbb{N}}$, $w = (1, \dots)$ or $w = (2, \dots)$. Then, by (105) we get

$$\sum_{l=1}^2 f \circ F_l(w) = \begin{cases} a_{11} + a_{21} & \text{se } w = (1, \dots) \\ a_{12} + a_{22} & \text{se } w = (2, \dots) \end{cases} \quad (107)$$

Now fix $a_{ij} = p_{ji}$, where p_{ij} are the entries of the row stochastic matrix P initially fixed. Then we get $a_{11} + a_{21} = p_{11} + p_{12} = 1$ e $a_{12} + a_{22} = p_{21} + p_{22} = 1$. Therefore for such choice of a_{ij} and for any $w \in I_m^{\mathbb{N}}$, the sum (107) equals 1. So, by (106), we get

$$\int \log\left(\sum_{i=1}^2 \frac{f \circ F_i}{f}\right) d\nu = - \sum_{i,j=1}^2 \pi_i p_{ij} \log p_{ij} = H(P) \quad (108)$$

Therefore,

$$\inf_{f \in \mathbb{B}^+} \int \log\left(\sum_{i=1}^2 \frac{f \circ F_i}{f}\right) d\nu \leq H(P) \quad (109)$$

◇

Now note that any positive function f can be written as

$$f(w) = \sum_{i,j=1}^2 a_{ij} p_{ji} 1_{\overline{ji}}(w)$$

Define

$$u(w) := \sum_{i,j=1}^2 a_{ij} 1_{\overline{ji}}(w)$$

and

$$g(w) := \sum_{i,j=1}^2 p_{ji} 1_{\overline{ji}}(w)$$

We have

$$\begin{aligned} \int_{I_2^{\mathbb{N}}} \log f d\nu &= \int_{I_2^{\mathbb{N}}} \sum_{i,j=1}^2 \log(a_{ij} p_{ji}) 1_{\overline{ji}} d\nu = \sum_{i,j=1}^2 \log(a_{ij} p_{ji}) \nu(\overline{ji}) \\ &= \sum_{i,j=1}^2 \pi_j p_{ji} \log(a_{ij} p_{ji}) = \sum_{i,j=1}^2 \pi_j p_{ji} \log(a_{ij}) + \sum_{i,j=1}^2 \pi_j p_{ji} \log(p_{ji}) \end{aligned} \quad (110)$$

If $w = (i, j, \dots)$, then $f \circ F_l(w) = a_{li} p_{il}$ and so

$$\sum_l f \circ F_l = \sum_l a_{li} p_{il}$$

We write

$$\mathcal{L}_g(u)(w) = \sum_l f \circ F_l(w) = \sum_l \sum_{i,j} a_{ij} p_{ji} 1_{ij}(F_l(w)) \quad (111)$$

We also have the following:

Lemma 5.

$$\int \mathcal{L}_g(\log u) d\nu = \int \log u d\nu \quad (112)$$

Proof We have

$$\int \log u d\nu = \int \sum_{i,j} \log(a_{ij}) 1_{ji} d\nu = \sum_{i,j} \log(a_{ij}) \nu(\overline{ji}) = \sum_{i,j} \log(a_{ij}) \pi_j p_{ji} \quad (113)$$

And also

$$\int \mathcal{L}_g(\log u) d\nu = \int \sum_l \sum_{i,j} \log(a_{ij}) p_{ji} 1_{\overline{ij}}(F_l(w)) d\nu$$

$$\begin{aligned}
 &= \sum_{i,j} \log(a_{ij}) p_{ji} \sum_l \int 1_{\overline{lj}}(F_l(w)) d\nu \\
 &= \sum_{i,j} \log(a_{ij}) p_{ji} \sum_l \nu(\overline{lj}) = \sum_{i,j} \log(a_{ij}) p_{ji} (\pi_1 p_{1j} + \pi_2 p_{2j}) = \sum_{i,j} \log(a_{ij}) \pi_j p_{ji}
 \end{aligned} \tag{114}$$

So,

$$\int \mathcal{L}_g(\log u) d\nu = \int \log u d\nu \tag{115}$$

□

Then, by using (110), (112) and (113),

$$\begin{aligned}
 &\int \log\left(\sum_{i=1}^k \frac{f \circ F_i}{f}\right) d\nu = \int \log\left(\sum_{l=1}^2 f \circ F_l\right) d\nu - \int \log f d\nu \\
 &= \int \log\left(\sum_{l=1}^2 f \circ F_l\right) d\nu - \left(\sum_{i,j=1}^2 \pi_j p_{ji} \log(a_{ij}) + \sum_{i,j=1}^2 \pi_i p_{ij} \log(p_{ij}) \right) \\
 &= \int \log(\mathcal{L}_g(u)) d\nu - \int \log u d\nu + H(P)
 \end{aligned} \tag{116}$$

$$= \int \log(\mathcal{L}_g(u)) d\nu - \int \mathcal{L}_g(\log u) d\nu + H(P) \tag{117}$$

We would like to show that

$$\int \log(\mathcal{L}_g(u)) d\nu - \int \mathcal{L}_g(\log u) d\nu \geq 0 \tag{118}$$

This follows immediately if we show that for $w = (i, j, \dots)$,

$$\log(\mathcal{L}_g(u))(w) \geq \mathcal{L}_g(\log u)(w) \tag{119}$$

The last expression follows from convexity. Indeed, to prove the above inequality, it is enough to show that for any $w = (i, j, \dots)$, we have

$$\log\left(\sum_l a_{li} p_{il}\right) \geq \sum_l p_{il} \log a_{li} \tag{120}$$

And such inequality is true, because the p_{il} are positive numbers with $\sum_l p_{il} = 1$, for any i , and the function \log is concave.

Therefore we conclude from (117) and (118) that

$$\int \log\left(\sum_{i=1}^k \frac{f \circ F_i}{f}\right) d\nu \geq H(P) \tag{121}$$

Conclusion By (109) and (121) we conclude that if ν is a Markov measure associated to a stochastic matrix P , then

$$\inf_{f \in \mathbb{B}^+} \int \log \left(\sum_{i=1}^2 \frac{f \circ F_i}{f} \right) d\nu = H(P), \quad (122)$$

and the function f such that

$$f(x) = \sum_{i,j=1}^2 p_{ij} 1_{\bar{i}\bar{j}}(x) \quad (123)$$

realizes the infimum.

◇

We conclude this section by stating the variational problem of pressure for our setting. We consider the the set of V_i , $i = 1, 2, \dots, k$ fixed, and we consider a variable set of W_i , $i = 1, 2, \dots, k$. In the normalized case, the different possible choices of p_i , $i = 1, 2, \dots, k$, (which means different choices of W_i , $i = 1, 2, \dots, k$) play here the role of the different Jacobians of possible invariant probabilities (see [15] II.1, and [12]) in Thermodynamic Formalism. In some sense the probabilities μ can be identified with the Jacobians (this is true at least for Gibbs probabilities of Hölder potentials [17]). The set of Gibbs probabilities for Hölder potentials is dense in the set of invariant probabilities [11].

Let $H : \mathcal{M}_N \rightarrow \mathcal{M}_N$ be a hermitian operator. We have the following problem. Define $F_0 : \mathcal{M}_F \rightarrow \mathbb{R}$,

$$F_0(\mu) := h_0(\mu) - \frac{1}{T} \text{tr}(H\rho_\mu) = \inf_{f \in \mathbb{B}^+} \int \log \left(\sum_{i=1}^k \frac{f \circ F_i}{f} \right) d\mu - \frac{1}{T} \text{tr}(H\rho_\mu),$$

where ρ_μ is the barycenter of μ , that is, the unique $\rho \in \mathcal{M}_N$ such that

$$l(\rho) = \int_{\mathcal{M}_N} l d\mu,$$

for all $l \in V^*$. Then, in order to find the associated Gibbs state we have to find $\hat{\mu} \in \mathcal{M}_F$ such that

$$F_0(\hat{\mu}) = \sup_{\mu \in \mathcal{M}_F} F_0(\mu).$$

We consider above each μ which is associated to a possible set of W_i .

◇

References

- [1] Baraviera, A., Lardizabal, C. F., A. Lopes, A. O., Terra Cunha, M. A dynamical point of view of Quantum Information: entropy, pressure and Wigner measures. Preprint Arxiv (2009).
- [2] Baraviera, A., Lardizabal, C. F., A. Lopes, A. O., Terra Cunha, M. A Thermodynamic Formalism for density matrices in Quantum Information. Preprint Arxiv (2009).
- [3] Benatti, F. Dynamics, Information and Complexity in Quantum Systems. Springer Verlag, 2009.
- [4] Benenti, G., Casati, G. Strini, G. Principles of Quantum Computation and Information, Vols. I and II, World Scientific, 2007.
- [5] Castro, G., Lopes, A. O. KMS States, Entropy and a Variational Principle for Pressure, to appear in Real Analysis Exchange (2009).
- [6] Gardiner, C. W., Zoller, P. Quantum Noise, Springer Verlag, 2004.
- [7] Gudder, S. Quantum Probability. Academic Press, 1988.
- [8] Gudder, S., Schindler, C. Quasi-discrete quantum Markov processes. Annales de l'Institut Henri Poincaré. Vol 56, no. 2, 1992.
- [9] Gudder, S. Quantum measure and Integration Theory, preprint (2009).
- [10] Lardizabal, C. F. Processos Estocásticos Quânticos, Tese de doutorado Prog. Posgrad. Mat. UFRGS - to appear (2010).
- [11] Lopes, A. O. Entropy and Large Deviation. NonLinearity Vol. 3, N. 2, 527-546, 1990.
- [12] Lopes, A. An analogy of the charge distribution on Julia sets with the Brownian motion. J. Math. Phys. 30 (9), 1989.
- [13] Lopes, A., Oliveira, E. Entropy and variational principles for holonomic probabilities of IFS. Discrete and Continuous Dynamical Systems Vol. 23, N, 3, 937-955 (2009) Series A.
- [14] Lozinski, A., Życzkowski, K., Słomczyński, W. Quantum iterated function systems, Physical Review E, Volume 68, 04610, 2003.
- [15] Mañé, R. Ergodic Theory, Springer Verlag, 1986.
- [16] Nielsen, M., Chuang, I. Quantum computation and quantum information. Cambridge University Press, 2000.
- [17] Parry, W., Pollicott, M. Zeta Functions and the Periodic Orbit Structure of Hyperbolic Dynamics. Société Mathématique de France. 187-188, Astérisque, 1990.
- [18] Słomczyński, W., Życzkowski, K. Quantum Chaos: an entropy approach. J. Math. Physics, 32 (1), 1994, p. 5674-5700.
- [19] Słomczyński, W. Dynamical Entropy, Markov Operators and Iterated Function Systems. Jagiellonian University Press, 2003.
- [20] Srinivas, M. D. Foundations of a quantum probability theory. Journal of Math. Phys., Vol. 16, No. 8, 1975.